

Tutorial 3, Statistical Mechanics: Concepts and applications 2016/17 ICFP Master (first year)

Maurizio Fagotti, Olga Petrova, Werner Krauth
Tutorial exercises

I. WORKSHEET: STATISTICAL MECHANICS AND THERMODYNAMICS

1. **Two level system.** [Source: Problem 6.3 from *Statistical Mechanics 2nd edition by Kerson Huang, 1987*]

Reminder: Entropy is given by

$$S = k \ln \Omega \quad (1)$$

where Ω is the number of microscopic configurations (in the quantum case, number of states) that correspond to a thermodynamic system in a state specified by certain macroscopic variables.

The inverse of the temperature is defined as

$$\frac{1}{T} = \left(\frac{\partial S}{\partial E} \right)_{V,N} \quad (2)$$

Consider a system of N free particles in which the energy of each particle can assume two and only two distinct values 0 and ϵ ($\epsilon > 0$). Denote by n_0 and n_1 , the occupation numbers of the energy level 0 and ϵ , respectively. The total energy of the system is E .

HINT # 1: Use ϵ as the unit of energy (i.e., set ϵ to 1) to simplify your expressions.

HINT # 2: You may find Stirling's approximation useful:

$$\ln x! \approx x \ln x - x$$

- (a) Find the entropy of such a system.
- (b) Find the temperature as a function of E , and show that it can be negative. What happens when a system of negative temperature is allowed to exchange energy with a system of positive temperature?

2. **Pauling residual entropy.** [Source: Pauling, *J. Am. Chem. Soc.* **12** (2680-2684), 1935.]




FIG. 1. (A) Local arrangement in water ice, showing oxygen ions (large white circles) and hydrogen ions (small black circles). (B) The corresponding picture for spin ice. (C) Pyrochlore lattice of corner-sharing tetrahedra, as occupied by the magnetic rare-earth ions in the spin ice materials $\text{Ho}_2\text{Ti}_2\text{O}_7$ and $\text{Dy}_2\text{Ti}_2\text{O}_7$. Source: Bramwell, Gingras, *Science* **294** (1495-1501), 2001.

Water molecules in ice are arranged in such a way that for every oxygen atom, there are two hydrogen atoms nearby, and two further away from it, as shown in Fig. 1(A). In 1935 Pauling used this to estimate the value of **residual entropy** - a nonzero value of entropy that systems, which can take multiple configurations at or near absolute zero, are said to have.

- (a) Estimate the value of residual entropy for an ice block containing N water molecules.

This problem came to a new light in 1999, when Ramirez and collaborators found that a rare-earth magnetic compound, $\text{Dy}_2\text{Ti}_2\text{O}_7$, reached the same value of residual entropy as water ice as the temperature tended towards zero. The source of the ground state degeneracy comes from the similar so-called ice-rule, which states that for every tetrahedron, making up the pyrochlore lattice [Fig. 1(C)] that Dy ions live on, there are two magnetic moments pointing in, and two pointing out [Fig. 1(B)].

3. Pressure.

Reminder: There is an important relation between the thermodynamic energy \bar{E} and the Hamiltonian $E(\{q\}, \{p\}, \lambda)$:

$$\overline{\frac{\partial E(\{q\}, \{p\}, \lambda)}{\partial \lambda}} = \left. \frac{\partial \bar{E}}{\partial \lambda} \right|_S. \quad (3)$$

Consider an external force acting on the body. Let the external parameter λ in Eq. (3) be \vec{r} , the radius of some surface element $d\vec{s}$. The force acting on that surface element is then given by

$$\vec{F} = \frac{\partial E(\{q\}, \{p\}, \vec{r})}{\partial \vec{r}}. \quad (4)$$

- (a) Find the mean force on a surface element. The magnitude of the force per unit area is called the **pressure**.

Hint: Change in volume is given by $d\vec{s} \cdot \vec{r}$

- (b) Consider two parts, in contact with one another, of a closed system in equilibrium. What can you say about the pressures of the two parts, P_1 and P_2 ?

4. Maxwell distribution.

- (a) Derive the probability distribution for the momentum of a non interacting particle.
- (b) What is the mean kinetic energy of the particle?
- (c) Find the mean square fluctuation of the velocity.