

Directed polymers in a random medium: $1/d$ expansion and the n -tree approximation

J Cook^{†‡} and B Derrida[†]

[†] Service de Physique Théorique§ de Saclay, 91191 Gif-sur-Yvette Cedex, France

[‡] Department of Physics, Edinburgh University, Edinburgh EH9 3JZ, UK

Received 25 October 1989

Abstract. We develop a method which allows one to obtain $1/d$ expansions for the problem of directed polymers in a random medium. Using this method, we expand thermal properties (such as the free energy, the specific heat), the overlaps and the fluctuations of the transverse displacement. Our results are consistent with the existence of a finite upper critical dimension above which the low-temperature phase is mean field like, i.e. characterised by a broad distribution of overlaps (broken replica symmetry) and transverse fluctuations which scale as the length of the polymer.

1. Introduction

The problem of directed polymers in a random medium can be formulated in its simplest lattice version as follows (Kardar 1987, Zhang 1987, Kardar and Zhang 1987, Cook and Derrida 1989b). For each bond, ij , of some regular lattice one chooses a random energy, ε_{ij} , according to a given probability distribution $\rho(\varepsilon_{ij})$. One then considers all directed walks of length L starting at a fixed origin on the lattice. A directed walk is, by definition, one stretched along a single longitudinal direction, with fluctuations in the transverse directions only. In this paper we shall consider walks on a d -dimensional hypercubic lattice, directed along the $(1, 1, \dots, 1)$ direction of the lattice. This means that at each step one coordinate must increase by one, so that if the walks commence at $\mathbf{r} = (x_1, \dots, x_d)$ they can end at any point $\mathbf{r}' = (x_1 + n_1, \dots, x_d + n_d)$ with $n_1 + \dots + n_d = L$. The energy, E_w , of a walk, w , is defined as

$$E_w = \sum_{ij \in w} \varepsilon_{ij} \quad (1)$$

where the summation runs over all the bonds, ij , visited by the walk w .

Three kinds of properties can be studied for such a system: thermal properties, overlaps and geometrical properties.

(i) *Thermal properties.* These can be obtained from the partition function Z :

$$Z_L(\mathbf{r}) = \sum_w \exp(-E_w/T). \quad (2)$$

In (2) the sum includes all the directed walks of length L starting from the point \mathbf{r} , and T is the temperature. From the knowledge of Z , one can calculate the free energy, the energy and the specific heat as functions of the temperature and then study the

§ Laboratoire de l'Institut de Recherche Fondamentale du Commissariat à l'énergie Atomique.

phase diagram. In the thermodynamic limit ($L \rightarrow \infty$), one can observe phase transitions between a high-temperature phase where the effect of disorder is weak and a low-temperature phase very similar to a spin-glass phase.

At zero temperature, the problem becomes an optimisation problem: the problem of finding the walk of lowest energy. In addition to the value of the ground-state energy, one can study its fluctuations (Zhang 1987, Kardar and Zhang 1987, Huse *et al* 1985) which are characterised by an exponent ω which depends on the dimension of the lattice

$$\langle E_{GS}^2 \rangle - \langle E_{GS} \rangle^2 \sim L^{2\omega} \quad (3)$$

where $\langle \rangle$ means an average over disorder.

(ii) *Overlaps*. As one can think of the low-temperature phase as a phase where the directed walks get trapped in deep valleys, it is useful, in order to study the statistical properties of the landscape, to define the overlap between two walks and the probability distribution of these overlaps (Derrida and Spohn 1988, Cook and Derrida 1989b). This overlap is defined by the fraction of their length that the two walks spend together.

(iii) *Geometrical properties*. The effect of disorder is to influence the transverse fluctuations of the directed walks. These can be characterised by an exponent ν :

$$Z^{-1} \sum_w (\mathbf{R}(w))^2 \exp(-E_w/T) \sim L^{2\nu} \quad (4)$$

where $\mathbf{R}(w)$ is the projection of the walk on the transverse directions.

The problem of directed polymers in a random medium has close analogies with two other problems in statistical mechanics: growth models and spin glasses.

The link with models of growing surfaces (Eden model or ballistic deposition) has been discussed by Kardar and Zhang (1987) and McKane and Moore (1988). They showed that the Kardar-Parisi-Zhang equation (Kardar *et al* 1986) for growth models could be interpreted as a continuous version of the directed polymer problem. This mapping allows one to relate the exponents ω and ν defined by (3) and (4) to the critical exponents characterising the fluctuations and roughness of the growing interface (Krug and Spohn 1988, Wolf and Kertész 1987, Kim and Kosterlitz 1989).

The link with spin glasses has been made in the mean-field limit (Derrida and Spohn 1988). It was shown in the mean-field theory of the directed polymer problem that the low-temperature phase is characterised by a broad distribution of overlaps as in Parisi's mean-field theory of spin glasses (Parisi 1980a, b, Mezard *et al* 1987). For spin glasses, the relevance of the Parisi solution in finite dimension has been a topic of controversy for the past decade (Sourlas 1986, Huse and Fisher 1987, 1988). One difficulty in trying to extend the Parisi solution to finite-dimensional systems is that the mean-field free energy is sufficiently complicated to make expansions around it technically very difficult (Bray and Moore 1979, de Dominicis and Kondor 1985). As the analytic expression for the free energy of the low-temperature phase is much simpler in the polymer problem, it should be an easier system in which to test the validity of the Parisi approach in finite-dimensional systems (Cook and Derrida 1989a).

Let us now summarise what is already known about the problem of directed polymers in a random medium.

A phase transition is known to exist for $d > 2+1$ (two transverse plus one longitudinal directions) (Imbrie and Spencer 1988). Above the transition temperature, T_c , the thermal properties take simple forms, the annealed and quenched free energies being the same. In the mean-field limit (Derrida and Spohn 1988) one can calculate

the transition temperature, whereas for finite-dimensional systems only bounds have been obtained (Cook and Derrida 1989b).

A lot of work has been done on determining the exponents ω and ν (see (3), (4)) at low temperature. In 1+1 dimensions the problem can be mapped on to a Burgers equation and a renormalisation analysis gives $\omega = \frac{1}{3}$ and $\nu = \frac{2}{3}$ (Forster *et al* 1977, Huse *et al* 1985). Little is known analytically about the exponents in higher dimension. A recent renormalisation group calculation (Halpin-Healy 1989, Nattermann and Renz 1988) has indicated a finite upper critical dimension $d_c = 4 + 1$. However, numerical simulations carried out by evaluating the surface scaling exponents for growth models have led to different conjectures. Wolf and Kertész (1987) suggest $\omega = 1/(2d - 1)$ and $\nu = d/(2d - 1)$ whilst Kim and Kosterlitz (1989) propose $\omega = 1/(d + 1)$ and $\nu = (d + 2)/2(d + 1)$ both giving no finite upper critical dimension. Recent simulations by Renz (1989) are intermediate between these two conjectures.

It is currently not established whether the exponents ω and ν vary continuously with dimension up to $d = \infty$ or whether they take their mean-field values, $\omega = 0$ and $\nu = \frac{1}{2}$, above a certain finite dimension d_c . In all cases, one expects these exponents to be related by a simple relation $\omega + 1 = 2\nu$ (Krug 1987).

Having summarised the problem and some of the results known already, let us now outline the rest of the paper and the new results we shall present. Firstly we shall discuss the mean-field version of the problem, recalling the methods of solution and the main results. Then, in section 3 we present the $1/d$ expansion method, first introduced in a previous work (Cook and Derrida 1989a), which is based on a series of approximations which we call ' n -tree' approximations. This will allow us to determine the $1/d$ expansion of the free energy and of the ground-state energy when the distribution $\rho(\varepsilon)$ of energies is exponential. We extend the method to obtain the fluctuations of the transverse displacement in section 4 and we calculate the overlap in section 5. In section 6 we discuss how the results would be modified by considering other distributions of disorder. Lastly in section 7 we test our expansion method (based on the n -tree approximation) on several problems which can be solved by alternative approaches.

2. The tree problem

A mean-field version of the problem of directed polymers in a random medium can be solved exactly (Derrida and Spohn 1988), for any distribution of disorder, by taking the regular lattice to be a branch of a tree (see figure 1). This problem can be solved by several methods: an analogy with travelling waves, a replica approach or by using results known on the GREM (Derrida and Gardner 1986). In this section we shall outline the travelling wave approach and present the main mean-field results. The replica approach is described in appendix 1.




Figure 1. Directed walk on a tree with $K = 2$.

Consider the branch of a tree, with coordination number, $K + 1$ (figure 1). For each bond ij of the lattice one chooses a random energy, ε_{ij} , according to a given distribution $\rho(\varepsilon_{ij})$. If $Z_L(0)$ is the partition function of the K^L directed walks of length L starting at 0, defined by (1) and (2), one can write a recursion relation connecting Z_L and Z_{L+1} :

$$Z_{L+1}(0) = \sum_{i=1}^K \exp(-\varepsilon_{0i}/T) Z_L(i) \quad (5)$$

where $Z_L(i)$ is the partition function of the walks in the i th branch and ε_{0i} is the energy of the bond connecting the origin to the i th branch. The energies ε_{ij} are independent random variables and so the partition function itself is random. It is therefore natural to consider a generating function of Z_L , $H_L(x)$ defined by

$$H_L(x) = \langle \exp[-\exp(-x/T) Z_L] \rangle. \quad (6)$$

Using $H_L(x)$ the recursion (5) takes the simple form

$$H_{L+1}(x) = \left(\int \rho(\varepsilon) H_L(x + \varepsilon) d\varepsilon \right)^K \quad (7)$$

where we have used the fact that $\varepsilon_{01}, \dots, \varepsilon_{0K}$ and $Z_L(1), \dots, Z_L(K)$ are independent random variables. Notice that (7) is independent of temperature. The temperature dependence only remains in the initial condition (8).

$$H_0(x) = \exp[-\exp(-x/T)]. \quad (8)$$

It can easily be seen from (6) and (8) that

$$H_L(x) \rightarrow \begin{cases} 1 & \text{when } x \rightarrow \infty \\ 0 & \text{when } x \rightarrow -\infty. \end{cases} \quad (9)$$

Hence $H_L(x)$ has the shape of a wavefront. Equation (7) is a travelling wave equation, belonging to a class of nonlinear equations of the diffusion-reaction type, and has properties similar to those of the KPP (Kolmogorov-Petrovsky-Piscounov) equation (Bramson 1983)

$$\frac{\partial H}{\partial L} = \frac{\partial^2 H}{\partial x^2} + H^2 - H. \quad (10)$$

Finding, for large L , the solution of the travelling wave equation (7) or (10) is not simple. However, it has been shown by Hammersley (1974) for (7) and by Bramson (1983) for (10), that for large L , the solutions of these equations become travelling waves, W , which move with velocity v

$$H_L(x) = W(x - vL - c(L)) \quad (11)$$

where $c(L)/L \rightarrow 0$ as $L \rightarrow \infty$.

In general the exact form of W is not known but the analytic expression of the velocity is easy to obtain. The velocity of the wave, v , depends upon the initial condition. For an initial condition of the form

$$H_0(x) = \begin{cases} 1 - e^{-\gamma x} & x \rightarrow \infty \\ 0 & x \rightarrow -\infty \end{cases} \quad (12)$$

with $H_0(x)$ a monotonic function, it has been shown (Hammersley 1974, Bramson 1983) that the velocity is given by

$$v = \begin{cases} G_1(\gamma) & \text{if } \gamma \leq \gamma_{\min} \\ G_1(\gamma_{\min}) = v_{\min} & \text{if } \gamma \geq \gamma_{\min} \end{cases}$$

where

$$G_1(\gamma) = \frac{1}{\gamma} \log \left(K \int \rho(\varepsilon) e^{-\gamma \varepsilon} d\varepsilon \right) \quad (13)$$

and γ_{\min} is the value of γ that minimises the velocity, i.e.

$$\frac{d}{d\gamma} G_1(\gamma)|_{\gamma_{\min}} = 0 \quad (14)$$

($G_1(\gamma)$ has in general a single minimum: this could be shown by using the fact that $\gamma G_1(\gamma)$ is always a convex function of γ). From (13) we see that the exponential decay, γ , of the initial condition controls the velocity of the travelling waves. If $\gamma < \gamma_{\min}$ the velocity varies continuously with γ , whereas if $\gamma \geq \gamma_{\min}$ the velocity remains fixed at its minimal value, v_{\min} .

These properties of the travelling waves ((11)–(14)) can be mapped onto properties of the partition function. Because of (6), the distribution of $\log Z_L$ is concentrated near the place where the wavefront is located. This implies from (11) that $-v$ can be identified with the free energy per unit length of the system, so we have

$$\lim_{L \rightarrow \infty} \frac{T}{L} \langle \log Z_L \rangle = \begin{cases} G_1(1/T) & T \geq T_c \\ G_1(\gamma_{\min}) & T \leq T_c \end{cases} \quad (15)$$

where

$$T_c \gamma_{\min} = 1. \quad (16)$$

This result is the same as that obtained by the replica method (equation (A1.12) of appendix 1). From (15) one sees that the entropy vanishes in the whole of the low-temperature phase, so that the system is completely frozen. One can conclude from (11) that the width of the probability distribution of $\log Z_L$ must be of order one, and this means that the energy fluctuation exponent, defined by (3), $\omega = 0$. It is not, however, possible to obtain an explicit expression for the ground-state energy fluctuations as this would require the knowledge of the shape of the travelling wave.

Another set of quantities which are of interest in this problem are the overlaps. If one defines $q_2(w, w')$ to be the fraction of their length that two walks w and w' , of length L , spend together, one can define the average overlap, $\langle q_2 \rangle$, to be

$$\langle q_2 \rangle = \left\langle \frac{1}{Z_L^2} \sum_w \sum_{w'} q_2(w, w') \exp \left(-\frac{E_w}{T} - \frac{E_{w'}}{T} \right) \right\rangle. \quad (17)$$

Similarly one can define the average overlap between m walks, $\langle q_m \rangle$. These overlaps are analogous to the overlaps which occur in the mean-field spin glass (Parisi 1983, Mezard *et al* 1984). Often it is useful to consider the probability distribution of the overlaps, $P_m(q)$. When replica symmetry breaking occurs one expects a many-valley landscape, in which the overlaps, q_m , will not be self-averaging. $P_2(q)$ will then be a non-trivial distribution function, which we define as

$$P_2(q) = \left\langle \frac{1}{Z_L^2} \sum_w \sum_{w'} \delta(q - q_2(w, w')) \exp \left(-\frac{E_w}{T} - \frac{E_{w'}}{T} \right) \right\rangle \quad (18)$$

with similar formulae pertaining for $P_m(q)$. In appendix 1, a replica approach is used to show, along with results (13)–(16), that the average overlap, $\langle q_2 \rangle$, and the distribution $P_2(q)$ are given by

$$\langle q_2 \rangle = \begin{cases} 1 - T/T_c & T \leq T_c \\ 0 & T \geq T_c \end{cases} \quad (19)$$

and

$$P_2(q) = \begin{cases} \frac{T}{T_c} \delta(q) + \left(1 - \frac{T}{T_c}\right) \delta(q-1) & T \leq T_c \\ \delta(q) & T \geq T_c \end{cases} \quad (20)$$

for the tree problem (Derrida and Spohn 1988).

The fact that $P_2(q)$ is not a single delta function for $T < T_c$ indicates a free energy landscape with a many-valley structure. Two walks lying in the same valley have an overlap $q_2 = 1$, whereas walks lying in different valleys have zero overlap.

To summarise, we have seen in this section that a travelling wave approach can be used to solve the mean-field problem of polymers in random media. The main results (13)–(16), (19), (20) for the mean-field limit will provide the basis for the $1/d$ expansions which we shall now present in sections 3, 4 and 5.

3. The $1/d$ expansion and the n -tree approximation

In this section we present the method of ' n -tree' approximations and show how this can be used to develop a $1/d$ expansion for the free energy of the directed polymer on a hypercubic lattice (Cook and Derrida 1989a). We use the expansion to derive formulae for the free energy, ground-state energy and specific heat in the low-temperature phase (equations (42), (46) and (44), respectively) and the transition temperature (equation (45)).

Consider the problem of directed polymers on a hypercubic lattice of dimension d . As usual one chooses a random energy, ε_{ij} , according to a given probability distribution, $\rho(\varepsilon)$, for each bond ij of the lattice. In this section and in sections 4 and 5 we shall choose (Dhar 1988)

$$\rho(\varepsilon) = \exp(-\varepsilon) \quad \varepsilon \geq 0. \quad (21)$$

This choice will make the calculation of several integrals we shall require much simpler. The use of other distributions will be discussed in section 6. Consider walks of length L , directed so that at each step one coordinate increases by one. This means that one directs the walks along the $(1, 1, \dots, 1)$ direction, so that if the walks commence at $\mathbf{r} = (x_1, \dots, x_d)$ they can end at any point $\mathbf{r}' = (x_1 + n_1, \dots, x_d + n_d)$ with $n_1 + \dots + n_d = L$. There are d^L directed walks of length L emanating from any given origin. The partition function, $Z_L(\mathbf{r})$, is defined by (1) and (2) as before.

One can write the following recursion for $Z_L(\mathbf{r})$:

$$Z_{L+1}(\mathbf{r}) = \sum_{i=1}^d a_i(\mathbf{r}) Z_L(\mathbf{r} + \mathbf{e}_i) \quad (22a)$$

with

$$a_i(\mathbf{r}) = \exp(-\varepsilon_{\mathbf{r}, \mathbf{r} + \mathbf{e}_i} / T) \quad (22b)$$

where \mathbf{e}_i are the unit vectors of the lattice in the directions in which a step is allowed, and the initial condition is

$$Z_0(\mathbf{r}) = 1. \quad (23)$$

The recursion (22a) is very similar to the recursion (5) which was valid for the tree. The main difference is that, in (22a), the $Z_L(\mathbf{r} + \mathbf{e}_i)$ are correlated. For example $Z_L(\mathbf{r} + \mathbf{e}_i)$ and $Z_L(\mathbf{r} + \mathbf{e}_j)$ are correlated because they both depend on $Z_{L-1}(\mathbf{r} + \mathbf{e}_i + \mathbf{e}_j)$.

To cope with these correlations in a systematic way we have developed a series of approximations which we call ' n -tree' approximations. These become exact in the limit $n \rightarrow \infty$. To obtain the n -tree approximation to the problem, one iterates (22a) n times exactly and then neglects the remaining correlations. In other words one takes account of the correlations on the first n steps of the lattice exactly and then constructs a tree from this motif, by neglecting other correlations. Diagrams of the 1-tree, 2-tree and 3-tree are shown for $d = 2$ in figure 2.




Figure 2. (a) 1-tree; (b) 2-tree; (c) 3-tree drawn for $d = 2$.

3.1. The 1-tree approximation

The first approximation, a 1-tree, neglects all correlations on the right-hand side of (22a). This gives the mean-field problem of (5), with $K = d$, the solution of which has been discussed in section 2. If we call the free energy per unit length of the n -tree $F_n(T)$, then from (13)–(16) we have

$$F_1(T) = \begin{cases} -G_1(1/T) & T \geq 1/\gamma_{\min} \\ -G_1(\gamma_{\min}) & T \leq 1/\gamma_{\min} \end{cases} \quad (24a)$$

$$T \leq 1/\gamma_{\min} \quad (24b)$$

where

$$G_1(\gamma) = \frac{1}{\gamma} \log(d \langle e^{-\gamma F} \rangle) \quad (24c)$$

and γ_{\min} is given by

$$\left. \frac{dG_1(\gamma)}{d\gamma} \right|_{\gamma_{\min}} = 0. \quad (24d)$$

We use, as before, $\langle \dots \rangle$ to denote an average over disorder, so here we have

$$\langle e^{-\gamma \epsilon} \rangle \equiv \int \rho(\epsilon) e^{-\gamma \epsilon} d\epsilon. \quad (25)$$

For the distribution (21) one sees that

$$G_1(\gamma) = \frac{1}{\gamma} \log \left(\frac{d}{\gamma + 1} \right). \quad (26)$$

If d is large one can expand the free energy given by (24) and (26) in powers of $1/d$, yielding

$$F_1(T) = \begin{cases} \frac{1}{ed} + \frac{1}{e^2 d^2} + \frac{3}{2e^3 d^3} + O(d^{-4}) & T \leq T_c \\ T \log[(1+T)/dT] & T \geq T_c \end{cases} \quad (27a)$$

$$(27b)$$

and one finds that the transition temperature, T_c , is given by

$$T_c = \frac{1}{\gamma_{\min}} = \frac{1}{ed} + \frac{2}{e^2 d^2} + \frac{9}{2e^3 d^3} + O(d^{-4}). \quad (28)$$

3.2. The 2-tree approximation

In the 1-tree approximation we have neglected all correlations in (22a). To extend the theory beyond the mean-field limit one must consider the correlations neglected so far. To start to do this we move to the 2-tree approximation. The 2-tree approximation is obtained by iterating (22a) twice to give

$$Z_{L+2}(\mathbf{r}) = \sum_{i=1}^d a_i(\mathbf{r}) a_i(\mathbf{r} + \mathbf{e}_i) Z_L(\mathbf{r} + 2\mathbf{e}_i) + \sum_{i=1}^{d-1} \sum_{j>i}^d [a_i(\mathbf{r}) a_j(\mathbf{r} + \mathbf{e}_i) + a_j(\mathbf{r}) a_i(\mathbf{r} + \mathbf{e}_j)] Z_L(\mathbf{r} + \mathbf{e}_i + \mathbf{e}_j) \quad (29)$$

and now neglecting the correlations between the Z_L on the RHS of (29). Again one ends up with a tree (see figure 2(b)), but now it has $d + d(d-1)/2$ branches rather than d branches. The branches are of two types: d branches have an energy which is the sum of two random energies $\epsilon_i + \epsilon_j$, whereas the other $d(d-1)/2$ branches have an effective energy, ϵ_{eff} , given by

$$\epsilon_{\text{eff}} = -T \log [a_i(\mathbf{r}) a_j(\mathbf{r} + \mathbf{e}_i) + a_j(\mathbf{r}) a_i(\mathbf{r} + \mathbf{e}_j)]. \quad (30)$$

The effective energy on these $d(d-1)/2$ branches depends upon the temperature and represents the fact that these branches are composed of two paths.

As a tree structure still remains the 2-tree can be solved exactly as before (equation (24)). If one defines

$$G_2(\gamma) = \frac{1}{2\gamma} \log(d \langle \exp[-\gamma(\epsilon_1 + \epsilon_2)] \rangle + \frac{1}{2} d(d-1) \times \langle \{ \exp[-(\epsilon_1 + \epsilon_2)/T] + \exp[-(\epsilon_3 + \epsilon_4)/T] \}^{\gamma T} \rangle) \quad (31)$$

then the free energy per unit length of the 2-tree is given by

$$F_2(T) = \begin{cases} -G_2\left(\frac{1}{T}\right) & T \geq T_c \\ -G_2(\gamma_{\min}) & T \leq T_c \end{cases} \quad (32a)$$

with γ_{\min} the solution of

$$\left. \frac{dG_2(\gamma)}{d\gamma} \right|_{\gamma_{\min}} = 0. \quad (32c)$$

The extra factor $\frac{1}{2}$ in (31) appears because when one iterates (29) L times the polymer is of length $2L$, not L . One sees that γ_{\min} now depends upon the temperature, because the effective energy on $d(d-1)/2$ of the branches was temperature dependent. T_c is still given by (16), i.e. is the solution of $T_c = 1/\gamma_{\min}(T_{jie0c})$.

3.3. The 3-tree approximation

To get successively better approximations to the d -dimensional lattice one can consider in turn a 3-tree, 4-tree, etc. . . . For example for the 3-tree

$$G_3(\gamma) = \frac{1}{3\gamma} \log[d(e^{-\gamma e})^3 + d(d-1)\langle f_2(T)^{\gamma T} \rangle + \frac{1}{6}d(d-1)(d-2)\langle f_3(T)^{\gamma T} \rangle] \quad (33a)$$

where

$$f_2(T) = a_1 a_2 a_3 + a_4 a_5 a_3 + a_4 a_6 a_7 \quad (33b)$$

and

$$f_3(T) = a_1 a_2 a_7 + a_1 a_6 a_{10} + a_4 a_3 a_7 + a_4 a_8 a_{11} + a_5 a_9 a_{10} + a_5 a_{12} a_{11} \quad (33c)$$

where $a_i = \exp(-\varepsilon_i/T)$.

One can see that $f_2(T)$ and $f_3(T)$ are the partition functions associated with the directed walks linking the two marked points on diagrams 2 and 3 of figure 3 respectively. It is clear that as the number of energies that must be averaged over increases the problem becomes much harder. So, although the aim would be to proceed to successive n -tree approximations, the complexity of the calculations that would be involved makes this approach impossible. However, for large dimension the problem is tractable.




Figure 3. f_i is the sum of the weights of all the directed walks connecting the two marked points on the diagram i .

3.4. The 2-tree for large d

When we expanded the tree solution for large d , equations (27a), (28), one saw that T_c was of order $1/d$. The key to obtaining the $1/d$ expansion is to keep in mind that for large dimension T_c is small, of order $1/d$, and γ_{\min} is large, of order d .

It is the fact that the relevant temperatures are small $\sim 1/d$ which simplifies the situation. To see this, consider (31). One needs to calculate

$$\langle f_1(T)^{\gamma T} \rangle = \langle \{ \exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T] \}^{\gamma T} \rangle \quad (34)$$

corresponding to figure 3(a). As T is at most of order $1/d$ in the low-temperature phase, the leading contribution to (34) will result from the region in the space of $\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4$ where one of the two terms on the RHS of (34) is much larger than the other. One can rewrite (34) as

$$\begin{aligned} \langle f_1(T)^{\gamma T} \rangle &= 2\langle \exp(-\gamma\varepsilon) \rangle^2 + \langle \{ \exp[-(\varepsilon_1 + \varepsilon_2)/T] \\ &\quad + \exp[-(\varepsilon_3 + \varepsilon_4)/T] \}^{\gamma T} \rangle - 2\langle \exp(-\gamma\varepsilon) \rangle^2. \end{aligned} \quad (35)$$

Using (21) one sees that the first term on the RHS of (35) is $2(\gamma+1)^{-2}$, whilst the second is only of order γ^{-4} . For example, it is easy to show that at $T=0$ the second term is $-2(3\gamma+4)(\gamma+1)^{-2}(\gamma+2)^{-3}$. Using the rearrangement (35) one can recast $G_2(\gamma)$ in the form

$$\begin{aligned} G_2(\gamma) &= \frac{1}{2\gamma} \log[d^2 \langle \exp(-\gamma\varepsilon) \rangle^2 + \frac{1}{2}d(d-1) \\ &\quad \times \langle \{ \exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T] \}^{\gamma T} \rangle - 2\langle \exp(-\gamma\varepsilon) \rangle^2]. \end{aligned} \quad (36)$$

3.5. The 3-tree for large d

To see how the high-dimension simplification works in a more complex case, consider $G_3(\gamma)$ given in (33). This was complicated because one had to evaluate $\langle f_2(T)^{\gamma T} \rangle$ and $\langle f_3(T)^{\gamma T} \rangle$. However, when d is large these quantities simplify in the same way as $\langle f_1(T)^{\gamma T} \rangle$ in (35). Take first $\langle f_2(T)^{\gamma T} \rangle$ defined in (33b). As T is small this is dominated by cases in which one of the three terms in (33b) is much larger than the other two. The next largest contribution comes from the situations when either the first pair or the last pair are much larger than the other term. Smaller again is the contribution due to cases when the first and last terms are of similar size, much greater than the middle term, and smallest of all is the effect of having all three terms of the same magnitude. Thus one can write

$$\begin{aligned} \langle f_2(T)^{\gamma T} \rangle &= 3\langle a^{\gamma T} \rangle^3 + 2\langle a^{\gamma T} \rangle \{ \langle (a_1 a_2 + a_3 a_4)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^2 \} \\ &\quad + \{ \langle (a_1 a_2 a_3 + a_4 a_5 a_6)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^3 \} \\ &\quad + \{ \langle (a_1 a_2 a_3 + a_4 a_5 a_3 + a_4 a_6 a_7)^{\gamma T} \rangle \\ &\quad - [\langle (a_1 a_2 a_3 + a_4 a_5 a_6)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^3] \\ &\quad - 2\langle a^{\gamma T} \rangle [\langle (a_1 a_2 + a_3 a_4)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^2] - 3\langle a^{\gamma T} \rangle^3 \}. \end{aligned} \quad (37)$$

The leading term is now $3(\gamma+1)^{-3}$, using (21), the second term is of order γ^{-5} , the third term is of order γ^{-6} and the last of order γ^{-7} . Similarly one can rewrite

$\langle f_3(T)^{\gamma T} \rangle$ as

$$\begin{aligned} \langle f_3(T)^{\gamma T} \rangle = & 6\langle a^{\gamma T} \rangle^3 + 6\langle a^{\gamma T} \rangle \{ \langle (a_1 a_2 + a_3 a_4)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^2 \} \\ & + 9\{ \langle (a_1 a_2 a_3 + a_4 a_5 a_6)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^3 \} \\ & + 6\{ \langle (a_1 a_2 a_3 + a_4 a_5 a_3 + a_4 a_6 a_7)^{\gamma T} \rangle \\ & - [\langle (a_1 a_2 a_3 + a_4 a_5 a_6)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^3] \\ & - 2\langle a^{\gamma T} \rangle [\langle (a_1 a_2 + a_3 a_4)^{\gamma T} \rangle - 2\langle a^{\gamma T} \rangle^2] - 3\langle a^{\gamma T} \rangle^3 \} + O(\gamma^{-8}). \end{aligned} \quad (38)$$

In (38) the first term, of order γ^{-3} , results from cases when one of the six walks comprising $f_3(T)$ dominates (33c). The second and third terms (of order γ^{-5} and γ^{-6}) represent cases when two of the six terms are of similar size, much larger than the rest and when these two dominant walks have one or no bonds in common, respectively. The fourth term is the effect of having three of the terms much larger than the other three, with one of the three dominant walks sharing one bond with each of the other two dominant walks. Further contributions to $\langle f_3(T)^{\gamma T} \rangle$ are of order γ^{-8} or smaller.

Using the rearrangements of (37) and (38) one can rewrite $G_3(\gamma)$ as

$$\begin{aligned} G_3(\gamma) = & \frac{1}{3\gamma} \log [d^3 \langle \exp(-\gamma \varepsilon) \rangle^3 + d^2 (d-1) \langle \exp(-\gamma \varepsilon) \rangle \\ & \times (\langle \{ \exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T] \}^{\gamma T} \rangle - 2\langle \exp(-\gamma \varepsilon) \rangle^2) \\ & + \frac{1}{2} d (d-1) (3d-4) (\langle \exp[-(\varepsilon_1 + \varepsilon_2 + \varepsilon_3)/T] \\ & + \exp[-(\varepsilon_4 + \varepsilon_5 + \varepsilon_6)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma \varepsilon) \rangle^3) + O(\gamma^{-4})]. \end{aligned} \quad (39)$$

3.6. The n -tree approximation for large d

Comparing (36) and (39), for large dimension, each of the terms in the argument of the logarithm on the RHS of the equations can be identified with a diagram in figure 4, and this allows us to obtain an expression for $G_n(\gamma)$ valid for arbitrary n . The first term in (36) and (39) represents a single walk of two and three steps respectively and can be identified with figure 4(a). Its contribution is a product of the embedding count $A(a)$ times the energy factor $B(a)$ given in table 1. The next term in (36) and (39) corresponds to figure 4(b), in which two walks bifurcate and rejoin immediately afterwards. The contribution is again the product $A(b)B(b)$ of table 1. Notice that an extra factor of two appears when $n=3$ as the bifurcation can occur at the first or second step. These two diagrams constitute all the possible arrangements of walks




Figure 4. The first six diagrams connecting points n steps apart (for $n \geq 4$).

Table 1. The embedding counts, A , and energy factors, B , of the diagrams of figure 4 for the diagrammatic expansion.

Diagram	Embedding count $A(\text{diagram})$
a	d^n
b	$\frac{1}{2}(n-1)d^{n-1}(d-1)$
c	$\frac{1}{2}(n-2)d^{n-2}(d-1)(3d-4)$
d	$(n-2)d^{n-2}(d-1)^2$
e	$\frac{1}{6}(n-2)(n-3)d^{n-2}(d-1)^2$
f	$\frac{1}{2}(n-3)d^{n-3}(d-1)(13d^2-40d+33)$
Diagram	Energy factor $B(\text{diagram})$
a	$\langle \exp(-\gamma\epsilon) \rangle^n$
b	$\langle \exp(-\gamma\epsilon) \rangle^{n-2} [\langle \exp[-(\epsilon_1 + \epsilon_2)/T] + \exp[-(\epsilon_3 + \epsilon_4)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^2]$
c	$\langle \exp(-\gamma\epsilon) \rangle^{n-3} [\langle \exp[-(\epsilon_1 + \epsilon_2 + \epsilon_3)/T] + \exp[-(\epsilon_4 + \epsilon_5 + \epsilon_6)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^3]$
d	$\langle \exp(-\gamma\epsilon) \rangle^{n-3} \{ \langle \exp[-(\epsilon_1 + \epsilon_2 + \epsilon_3)/T] + \exp[-(\epsilon_4 + \epsilon_5 + \epsilon_6)/T] \rangle^{\gamma T} + \exp[-(\epsilon_4 + \epsilon_6 + \epsilon_7)/T] \rangle^{\gamma T} - [\langle \exp[-(\epsilon_1 + \epsilon_2 + \epsilon_3)/T] + \exp[-(\epsilon_4 + \epsilon_5 + \epsilon_6)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^3] - 2\langle \exp(-\gamma\epsilon) \rangle [\langle \exp[-(\epsilon_1 + \epsilon_2)/T] + \exp[-(\epsilon_3 + \epsilon_4)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^2] - 3\langle \exp(-\gamma\epsilon) \rangle^3 \}$
e	$\langle \exp(-\gamma\epsilon) \rangle^{n-4} [\langle \exp[-(\epsilon_1 + \epsilon_2)/T] + \exp[-(\epsilon_3 + \epsilon_4)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^2]^2$
f	$\langle \exp(-\gamma\epsilon) \rangle^{n-4} [\langle \exp[-(\epsilon_1 + \epsilon_2 + \epsilon_3 + \epsilon_4)/T] + \exp[-(\epsilon_5 + \epsilon_6 + \epsilon_7 + \epsilon_8)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^4]$

linking two points two steps apart. The third term in (39) can be identified with figure 4(c) in which two walks split and rejoin three steps later. Again its contribution is given by $A(c)B(c)$ in table 1.

Hence, to be able to evaluate $G_n(\gamma)$ one need only consider all the different ways of linking two points n steps apart, starting by utilising as few bonds as possible and gradually increasing the number of bonds used by the walks to obtain higher orders in the expansion. For each diagram one has to determine the number of ways of embedding it in the lattice, $A(\text{diagram})$, and the corresponding energy factor, $B(\text{diagram})$. The diagrams of figure 4 have previously been used to calculate the high-dimension expansion of the directed bond percolation threshold (Blease 1977).

Using table 1, $G_n(\gamma)$ can be written as

$$\begin{aligned}
 G_n(\gamma) &= \frac{1}{n\gamma} \log[d^n \langle \exp(-\gamma\epsilon) \rangle^n \\
 &\quad + \frac{1}{2}(n-1)d^{n-1}(d-1) \langle \exp(-\gamma\epsilon) \rangle^{n-2} (\langle \exp[-(\epsilon_1 + \epsilon_2)/T] \\
 &\quad + \exp[-(\epsilon_3 + \epsilon_4)/T] \rangle^{\gamma T} - 2\langle \exp(-\gamma\epsilon) \rangle^2) \\
 &\quad + \frac{1}{2}(n-2)d^{n-2}(d-1)(3d-4) \langle \exp(-\gamma\epsilon) \rangle^{n-3} \\
 &\quad \times (\langle \exp[-(\epsilon_1 + \epsilon_2 + \epsilon_3)/T] + \exp[-(\epsilon_4 + \epsilon_5 + \epsilon_6)/T] \rangle^{\gamma T} \\
 &\quad - 2\langle \exp(-\gamma\epsilon) \rangle^3) + O(\gamma^{-4})] \\
 &= \frac{1}{n\gamma} \log \left(\sum_{\text{diagrams}} A(\text{diagram}) B(\text{diagram}) \right). \tag{40}
 \end{aligned}$$

In general a diagram will contribute at order γ^{-p} in the argument of the log in (40) if this diagram contains p extra bonds compared with figure 4(a). So to expand (40) to a given order in $1/\gamma = O(1/d)$, one needs to consider a finite number of diagrams. Having obtained $G_n(\gamma)$, the free energy per unit length then follows as before (e.g. (13)–(16))

$$F_n(T) = \begin{cases} -G_n(1/T) & T \geq T_c \\ -G_n(\gamma_{\min}) & T \leq T_c \end{cases} \quad (41a)$$

$$(41b)$$

with

$$\left. \frac{dG_n(\gamma)}{d\gamma} \right|_{\gamma_{\min}} = 0 \quad (41c)$$

and T_c given by (16).

Using the expansion for $G_n(\gamma)$ computed from table 1 (equation (40)) with the energy distribution (21), one can show that in the low-temperature phase

$$F_n(T) = \frac{1}{ed} + \frac{1}{e^2 d^2} + \frac{3}{2e^3 d^3} - \frac{1}{e^3 d^3} \left(\frac{n-1}{n} \right) [g(Ted) - 3] + O(d^{-4}) \quad (42a)$$

with

$$g(t) = t \int_0^\infty du (2+tu) [(1+e^{-u})^t - 1] \quad (42b)$$

and in the high-temperature phase (to all orders in $1/d$)

$$F_n(T) = T \log \left(\frac{T+1}{Td} \right) \quad T \geq T_c. \quad (43)$$

Notice (27a), (42a) that the first correction to the tree calculation for $T \leq T_c$ occurs at order d^{-3} , not d^{-2} , because the first corrective diagram includes two extra bonds. Also, one sees from (43) that in the high-temperature phase the quenched and annealed free energies are equal (Imbrie and Spencer 1988, Cook and Derrida 1989b). From (42a) the specific heat per unit length, $C_n(T)$, in the low-temperature phase can be shown to be

$$\begin{aligned} C_n(T) &= \frac{T}{ed} \left(\frac{n-1}{n} \right) g''(Ted) + O\left(\frac{1}{\gamma^2}\right) \\ &\simeq \frac{T}{ed} \left(\frac{n-1}{n} \right) \frac{\pi^2}{3} \quad \text{for } T \text{ small.} \end{aligned} \quad (44)$$

The low-temperature phase is no longer frozen as it was in the mean-field limit (Derrida and Spohn 1988). Instead one finds here a specific heat linear in temperature at low temperature. The transition temperature, T_c , can be calculated using (41c), (40) and (16) to yield

$$T_c = \frac{1}{ed} + \frac{2}{e^2 d^2} + \frac{1}{e^3 d^3} \left[\frac{9}{2} + \left(\frac{n-1}{n} \right) g'(1) \right] + O(d^{-4}) \quad (45)$$

where $g'(1) = 7.527\,827 \dots$ (see (42b)).

These results (42)–(45) were obtained using figures 4(a) and 4(b). One can go to higher orders in $1/d$ by including the next diagrams. For example, if one includes all the diagrams of figure 4, one can obtain for the ground-state energy per unit length

$$E_{\text{GS}} = \frac{1}{de} + \frac{1}{(de)^2} + \frac{1}{(de)^3} \left(\frac{9}{2} - \frac{3}{n} \right) + \frac{1}{(de)^4} \left(\frac{110}{3} - 3e + \frac{1}{n} (3e - 64) \right) \\ + \frac{1}{(de)^5} \left(\frac{11297}{24} - 74e + \frac{1}{n} (144e - 1369) + \frac{1}{n^2} \frac{81}{2} \right) + O\left(\frac{1}{d^6}\right). \quad (46)$$

(Notice that (46) is only valid for $n \geq 3$ because the embedding counts of table 1 are only valid for $n \geq 3$.)

The calculation of the ground-state energy in powers of $1/d$ is technically simpler than that of the free energy because all the energy factors B of table 1 can be calculated explicitly in the limit $T \rightarrow 0$ (table 2) for the exponential distribution (21).

Table 2. The energy factors, B , of table 1 calculated at $T = 0$ for the exponential distribution (21).

Diagram	Energy factor $B(\text{diagram})$ at $T = 0$
<i>a</i>	$\frac{1}{(\gamma + 1)^n}$
<i>b</i>	$\frac{-(6\gamma + 8)}{(\gamma + 1)^n (\gamma + 2)^3}$
<i>c</i>	$\frac{-(20\gamma^2 + 50\gamma + 32)}{(\gamma + 1)^n (\gamma + 2)^5}$
<i>d</i>	$\frac{71\gamma^4 + 555\gamma^3 + 1581\gamma^2 + 1937\gamma + 864}{(\gamma + 1)^n (\gamma + 2)^5 (\gamma + 3)^3}$
<i>e</i>	$\frac{(6\gamma + 8)^2}{(\gamma + 1)^n (\gamma + 2)^6}$
<i>f</i>	$\frac{-(70\gamma^3 + 252\gamma^2 + 308\gamma + 128)}{(\gamma + 1)^n (\gamma + 2)^7}$

All the results (41)–(46) obtained so far are $1/d$ expansions valid for the n -tree problems. As the n -tree problem is identical to the hypercubic lattice in the limit $n \rightarrow \infty$, we expect these results (41)–(46) to be valid for the hypercubic lattice in the limit $n \rightarrow \infty$.

So the $1/d$ expansion of the free energy, the specific heat and the ground-state energy for the hypercubic lattice are given by putting $n = \infty$ in (42), (44) and (46).

For these results to be correct, one needs the limits $n \rightarrow \infty$ and $d \rightarrow \infty$ to commute. We have not been able to prove that these two limits commute. However, in all the $1/d$ expansions presented in this paper, the coefficients are always simple rational functions of n which have a finite limit for $n \rightarrow \infty$. So there is no indication that the limit $n \rightarrow \infty$ is singular for large d . Moreover, we shall see in section 7 that there are some cases which can be solved without the n -tree approximation and for which the exchange of these limits $n \rightarrow \infty$ and $d \rightarrow \infty$ gives the right answer.

4. The fluctuations of the transverse displacement

In this section we calculate the fluctuations of the transverse displacement of the polymer. We shall see that one can define two types of transverse fluctuations: thermal and disorder fluctuations. Then the $1/d$ expansion, presented in the previous section, will be extended to include a chemical potential, enabling one to calculate the transverse fluctuations on a d -dimensional hypercubic lattice.

Let us first define what is meant by transverse fluctuations. Consider the problem of polymers on a d -dimensional hypercubic lattice, directed along the $(1, 1, \dots, 1)$ direction, as described in section 3. If a polymer of length L , emanating from the origin, ends at the point (x_1, \dots, x_d) , its transverse displacement is given by the vector \mathbf{R}

$$\mathbf{R} = \left(x_1 - \frac{L}{d}, \dots, x_d - \frac{L}{d} \right) = (R_1, \dots, R_d). \quad (47)$$

We want to consider the fluctuations in the quantities R_i ($i = 1, \dots, d$). These can be of two types: thermal fluctuations, given by

$$\langle \bar{R}_i^2 - \bar{R}_i^2 \rangle \quad (48a)$$

and disorder fluctuations, given by

$$\langle \bar{R}_i^2 \rangle - \langle \bar{R}_i \rangle^2 = \langle \bar{R}_i^2 \rangle \quad (48b)$$

since $\langle \bar{R}_i \rangle = 0$ for symmetry reasons.

We use the notation that $\langle \rangle$ denotes an average over disorder and $\bar{}$ a configurational or thermal average. Often the quantity $\sum_{i=1}^d \langle \bar{R}_i^2 \rangle$ is considered; this scales as the length of the polymer, L , to an exponent 2ν , i.e.

$$\sum_{i=1}^d \langle \bar{R}_i^2 \rangle = d \langle \bar{R}_i^2 \rangle \sim L^{2\nu}. \quad (49)$$

As was mentioned in section 1, little is known analytically about ν as a function of dimension except that $\nu = \frac{1}{2}$ in the mean-field limit and that $\nu = \frac{2}{3}$ in $d = 1 + 1$ dimensions.

4.1. The relation between thermal and disorder fluctuations

Let us now derive a relation between thermal and disorder fluctuations valid for any tree problem. Although the notion of transverse displacement has no *a priori* meaning in a tree problem, we can identify each direction $1 \leq i \leq d$ of a tree of d branches with one direction i on the hypercubic lattice and say that a walk on a tree has a transverse displacement R_i if it does $L/d + R_i$ steps along the i th branch.

Let us now attempt to evaluate the fluctuations (48) for the tree problem in the low-temperature phase. We have seen, (19) and (20), that in the low-temperature phase the overlaps are either 0 or 1. This implies that there are well defined valleys and that the thermal fluctuations inside a valley vanish. Therefore one can think of a landscape composed of a number of valleys. Each valley α is characterised by some transverse displacement $R_1 = c_\alpha$ in direction 1 and by a weight ω_α . Because the overlap between two walks inside the same valley is 1, all the walks inside the same valley α have the same transverse displacement c_α . Moreover, since the overlap between two different valleys is 0, the displacements c_α and c_β for two valleys are uncorrelated. Therefore one expects that the c_α satisfy

$$\langle c_\alpha \rangle = 0 \quad \langle c_\alpha c_\beta \rangle = \langle c_\alpha \rangle \langle c_\beta \rangle \quad \text{if } \alpha \neq \beta \quad (50)$$

where $\langle \rangle$ is the disorder average. Since the only thermal fluctuations are due to relative weights, ω_α , of the valleys, one has

$$\overline{R_1} = \sum_{\alpha} c_{\alpha} \omega_{\alpha} \quad (51a)$$

$$\overline{R_1^2} = \sum_{\alpha} c_{\alpha}^2 \omega_{\alpha}. \quad (51b)$$

To obtain the thermal and the disorder fluctuations, we now need to average over disorder. The weights ω_{α} satisfy

$$\sum_{\alpha} \omega_{\alpha} = 1 \quad (52a)$$

$$\left\langle \sum_{\alpha} \omega_{\alpha}^2 \right\rangle = 1 - T\gamma_{\min}. \quad (52b)$$

Equation (52a) expresses the fact that the weights of the valleys must add up to unity and (52b) is derived in appendix 1 (equation (A1.16)). If we consider that the c_{α} and the ω_{α} are not correlated, we get

$$\begin{aligned} \langle \overline{R_1} \rangle &= 0 \\ \langle \overline{R_1^2} \rangle &= \left\langle \sum_{\alpha} \omega_{\alpha} \right\rangle \langle c_{\alpha}^2 \rangle = \langle c_{\alpha}^2 \rangle \\ \langle \overline{R_1^2} \rangle &= \left\langle \sum_{\alpha} \omega_{\alpha}^2 \right\rangle \langle c_{\alpha}^2 \rangle = (1 - T\gamma_{\min}) \langle c_{\alpha}^2 \rangle \end{aligned} \quad (53)$$

which give for the thermal fluctuations

$$\langle \overline{R_1^2} \rangle - \langle \overline{R_1^2} \rangle = \langle c_{\alpha}^2 \rangle T\gamma_{\min} \quad T \leq T_c \quad (54)$$

and for the disorder fluctuations:

$$\langle \overline{R_1^2} \rangle = \langle c_{\alpha}^2 \rangle (1 - T\gamma_{\min}) \quad T \leq T_c. \quad (55)$$

The disorder and thermal fluctuations take on similar forms, their ratio being $(1 - T\gamma_{\min})/T\gamma_{\min}$. The disorder fluctuations vanish at T_c . This result is very similar to the fact that at T_c the quenched and annealed free energies become identical.

4.2. Calculation of the thermal fluctuations

We shall now study the transverse fluctuations on the hypercubic lattice using the $1/d$ expansion. This can be done by adding a chemical potential, μ , to count the number of steps in direction 1

$$Z_L(\mathbf{r}) = \sum_w \exp[-E_w/T + \mu(R_1(w) + L/d)] \quad (56)$$

where the sum runs over all walks of length L emanating from the point \mathbf{r} and $(R_1(w) + L/d)$ is the number of steps of walk w in direction 1.

The average number of steps taken in the direction marked by μ , direction 1, is then given by

$$\langle \overline{R_1} \rangle + \frac{L}{d} = \frac{d}{d\mu} \langle \log Z_L \rangle \Big|_{\mu=0} \quad (57)$$

and the thermal fluctuation of R_1 is given by the second derivative of $\langle \log Z_L \rangle$

$$\langle \overline{R_1^2} \rangle - \langle \overline{R_1} \rangle^2 = \frac{d^2}{d\mu^2} \langle \log Z_L \rangle \Big|_{\mu=0}. \quad (58)$$

The method of n -tree approximations and the resulting $1/d$ expansion can be used as in section 3 to calculate the free energy of problem (56), providing that care is taken to include an extra factor, $e^{\mu\gamma T}$, for each step taken in the marked direction. For example, the free energy per unit length of the 1-tree approximation, $F_1(T, \mu)$, can be obtained using (41) from $G_1(\gamma)$:

$$G_1(\gamma) = \frac{1}{\gamma} \log \left[\left(\frac{d-1+e^{\mu\gamma T}}{\gamma+1} \right) \right] \quad (59)$$

where the distribution (21) has again been used. The inclusion of the chemical potential has changed the embedding count of figure 4(a) from d^n to $(d-1+e^{\mu\gamma T})^n$. The embedding counts for the diagrams of figure 4, obtained using the chemical potential, are given in table 3. Combining the new embedding counts (table 3) with the energy factors (table 1), as in (40), it is then possible to find $G_n(\gamma)$ and hence the $1/d$ expansion for the free energy, $F_n(T, \mu)$.

From $F_n(T, \mu)$ one can calculate the thermal fluctuations on the n -tree (58) by

$$\langle \bar{R}_1^2 \rangle - \langle \bar{R}_1^2 \rangle = - \frac{L}{T} \frac{d^2}{d\mu^2} F_n(T, \mu) \Big|_{\mu=0}. \quad (60)$$

This formula is valid at all temperatures.

4.3. The disorder fluctuations in the low-temperature phase

From (54), (55) and (60) one can obtain the disorder fluctuations.

$$\langle \bar{R}_1^2 \rangle = - \frac{(1 - T\gamma_{\min})}{T^2 \gamma_{\min}} L \frac{d^2}{d\mu^2} F_n(T, \mu) \Big|_{\mu=0}. \quad (61)$$

Notice that (61) is only valid for a tree problem, since (54) and (55) only apply to this case.

Table 3. The embedding counts, A , for the diagrams of figure 4 when a chemical potential, μ , is included to count the number of steps in the marked direction.

Diagram	Embedding count
<i>a</i>	$(d-1+e^{\mu\gamma T})^n$
<i>b</i>	$\frac{1}{2}(n-1)(d-1)^{n-1} \left(1 + \frac{e^{\mu\gamma T}}{d-1}\right)^{n-2} (d-2+2e^{\mu\gamma T})$
<i>c</i>	$\frac{1}{2}(n-2)(d-1)^{n-2} \left(1 + \frac{e^{\mu\gamma T}}{d-1}\right)^{n-3} (3d^2-13d+14+(9d-16)e^{\mu\gamma T} + 2e^{2\mu\gamma T})$
<i>d</i>	$(n-2)(d-1)^{n-2} \left(1 + \frac{e^{\mu\gamma T}}{d-1}\right)^{n-3} (d^2-4d+4+(3d-5)e^{\mu\gamma T} + e^{2\mu\gamma T})$
<i>e</i>	$\frac{1}{8}(n-2)(n-3)(d-1)^{n-2} \left(1 + \frac{e^{\mu\gamma T}}{d-1}\right)^{n-4} (d-2+2e^{\mu\gamma T})^2$
<i>f</i>	$(n-3)(d-1)^{n-3} \left(1 + \frac{e^{\mu\gamma T}}{d-1}\right)^{n-4} \left[\frac{13}{2}d^3-46d^2+109d-86 + (26d^2-105d+107)e^{\mu\gamma T} + \left(\frac{25}{2}d-22\right)e^{2\mu\gamma T} + e^{3\mu\gamma T} \right]$

From (60), (61) one could obtain the complete temperature dependence of both the thermal and disorder fluctuations. However, as the calculations are greatly simplified at zero temperature we shall give the results in this limit only. At $T=0$ the thermal fluctuations vanish, whereas the disorder fluctuations are given by

$$\sum_{i=1}^d \langle \bar{R}_i^2 \rangle = L \left[1 - \frac{1}{d} + \frac{6}{e^2 d^3} \left(\frac{n-1}{n} \right) + \frac{(136-6e)n-276+6e}{ne^3 d^4} + \frac{(3384-376e)n+756e-10454+(108/n)}{ne^4 d^5} + O(d^{-6}) \right]. \quad (62)$$

Taking the limit $n \rightarrow \infty$ one gets a $1/d$ expansion for the disorder transverse fluctuation at zero temperature for the hypercubic lattice. As for the free energy, each term of the expansion has a regular n -dependence when $n \rightarrow \infty$ and so it is reasonable to assume that the limits $d \rightarrow \infty$ and $n \rightarrow \infty$ can be exchanged.

We see from (62) that the $1/d$ expansion does not change the value of ν , which remains $\frac{1}{2}$. This would be consistent with the fact that $\nu = \frac{1}{2}$ above a certain upper critical dimension, d_c , at which the coefficient of L (in the RHS of (62) and for $n \rightarrow \infty$) would diverge. The series (62) is, however, too short to make any reliable prediction for d_c . Moreover, it could happen that the $1/d$ series (62) (for $n = \infty$) has a zero radius of convergence, as is the case in some $1/d$ expansions (Fisher and Singh 1989).

The transverse fluctuations have an interesting temperature dependence. Let us just discuss now their behaviour to leading order in $1/d$. From (59), (60) and (61), one finds that the thermal fluctuations are given by,

$$\sum_{i=1}^d \langle \bar{R}_i^2 \rangle - \langle \bar{R}_i^2 \rangle = \begin{cases} L \left(1 - \frac{1}{d} \right) \frac{T}{T_c} + O\left(\frac{1}{d^2}\right) & \text{for } T \leq T_c \\ L \left(1 - \frac{1}{d} \right) & \text{for } T > T_c \end{cases} \quad (63)$$

and the disorder fluctuations are given by

$$\sum_{i=1}^d \langle \bar{R}_i^2 \rangle = \begin{cases} L \left(1 - \frac{1}{d} \right) \left(1 - \frac{T}{T_c} \right) + O\left(\frac{1}{d^2}\right) & \text{for } T \leq T_c \\ 0 & \text{for } T > T_c \end{cases} \quad (64)$$

(see figure 5).

The fact that we have two kinds of transverse fluctuations in this problem is reminiscent of the susceptibility of spin glasses for which, at low temperature, two susceptibilities can be defined (DC susceptibility and AC susceptibility, in the limit of zero frequency).

The inclusion of a chemical potential, as in (56), also enables one to consider the problem of polymers in a random medium in different geometries. For example, one can consider the problem in which the partition sum (2) runs over all walks with a fixed number of steps, N , in a given direction rather than over all walks of fixed length L . This problem is related to a problem considered by Dhar (1988), who showed that, on a d -dimensional hypercubic lattice, the ground-state energy for distribution (21) is given to leading order by $\log d/d$, for d large. Using our $1/d$ expansion method based on the n -tree approximation, it is possible to recover this leading mean-field term and calculate corrections to it.




Figure 5. The temperature dependence of disorder (full line) and thermal (broken line) transverse fluctuation, equations (63), (64).

5. The overlaps

As mentioned in section 1, the problem of directed polymers in random media has a close analogy with spin glasses. The mean-field spin glass (Parisi 1980a, b) and the mean-field polymer problem (Derrida and Spohn 1988) both have low-temperature phases characterised by broken replica symmetry. This is displayed in the polymer problem by the probability distribution of overlaps, $P_2(q)$ (see (17) and (18) for definition), being a sum of two delta functions (see appendix 1, equation (A1.14) and section 2). The question of replica symmetry breaking in the finite-dimensional spin glass remains controversial and for this reason it is interesting to examine the overlaps for the polymer problem in finite dimension. In this section, therefore, we shall study the overlaps on a d -dimensional hypercubic lattice by extending the n -tree approach, obtaining the probability distribution of the overlap of two walks (70), (71).

The n -tree approximations to the directed polymer problem, defined in section 3, retain a general tree structure. Hence, if one performs the replica calculation of appendix 1 on the n -tree one will see that, below T_c , $P_2(q)$ remains a sum of two delta functions with weights $T\gamma_{\min}$ and $1 - T\gamma_{\min}$. However, in the n -tree approximation some of the branches of the tree structure are composed of two or more paths. Consider what can occur when two walks pass down such a branch. The walks can both take the same path, giving an overlap of 1 for this section, or they can take different routes, giving an overlap less than 1 for this section. So, instead of having one delta function at $q_2 = 0$ and the other at $q_2 = 1$, the latter will be shifted to a lower value of q_2 .

To explain how one can calculate the position of the shifted delta function, let us consider as an example a polymer of length L in the 2-tree approximation. Here one has only two kinds of branch (see figure 2(b)): those which consist of only one path (type A), and so contribute $2/L$ to the overlap q_2 and those which are made up of two paths (type B), which contribute between $1/L$ (at $T = \infty$) and $2/L$ (at $T = 0$) to q_2 , depending on the temperature. To determine the distribution function of the overlaps, one therefore needs to know the fraction of each type of branch used by the walks and the contribution to the overlap from the type B branches, both as functions of temperature.

As is shown in appendix 2, the typical fraction of branches used in the walk that are of type B and have energies $\varepsilon_1 + \varepsilon_2$ and $\varepsilon_3 + \varepsilon_4$ on the constituent links is given, in

the low-temperature phase, by

$$\frac{1}{2}d(d-1)\{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}}\rho(\varepsilon_1)\rho(\varepsilon_2)\rho(\varepsilon_3)\rho(\varepsilon_4) \\ \times \exp[-2\gamma_{\min}G_2(\gamma_{\min})] \quad (65)$$

where $G_2(\gamma)$ is given by (36).

It now remains to calculate the contribution to the overlap, q_m , of such a branch. This is simply given by

$$\frac{\exp[-m(\varepsilon_1 + \varepsilon_2)/T] + \exp[-m(\varepsilon_3 + \varepsilon_4)/T]}{\{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^m} \quad (66)$$

as one has an overlap 1 if all the walks go through the same path and overlap 0 otherwise. Combining (65) and (66), one can conclude that $P_2(q)$ for the 2-tree consists of two delta functions, one at $q = 0$ with weight $T\gamma_{\min}$ and the other at

$$q_2 = (d\langle \exp(-\varepsilon\gamma_{\min}) \rangle^2 + \frac{1}{2}d(d-1)\{\{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}-2} \\ \times \{\exp[-2(\varepsilon_1 + \varepsilon_2)/T] + \exp[-2(\varepsilon_3 + \varepsilon_4)/T]\}\}) \\ \times \exp[-2\gamma_{\min}G_2(\gamma_{\min})]. \quad (67)$$

Having seen how the calculation proceeds for the 2-tree, it is easy to generalise to the n -tree. The relative contribution to q_m from each branch like figure 4(a) is always 1. The relative contribution from branches like figure 4(b) will be

$$1 - \frac{2}{n} + \frac{2}{n} \frac{\exp[-m(\varepsilon_1 + \varepsilon_2)/T] + \exp[-m(\varepsilon_3 + \varepsilon_4)/T]}{\{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^m}. \quad (68)$$

This is just the direct generalisation of (66): the walks have to be together for a fraction $1 - 2/n$ of the branch and can then take two paths for a fraction $2/n$ of the branch. So for the n -tree the average overlap $\langle q_2 \rangle$ takes the form

$$\langle q_2 \rangle = \left[1 - \left(\frac{n-1}{n} \right) \frac{(d-1)}{d} \langle \exp(-\varepsilon\gamma_{\min}) \rangle^{-2} \right. \\ \times (\langle \{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}} \rangle \\ - \langle \{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}-2} \rangle \\ \times \{\exp[-2(\varepsilon_1 + \varepsilon_2)/T] + \exp[-2(\varepsilon_3 + \varepsilon_4)/T]\} \rangle \\ \left. + O(d^{-3}) \right] (1 - T\gamma_{\min}) \quad (69)$$

where γ_{\min} is given by (40) and (41).

Evaluating the averages, using (21) and taking the limit $n \rightarrow \infty$ (again assuming the commutativity of limits) yields that the delta function at non-zero overlap in $P_2(q)$ is shifted from $q_2 = 1$ to

$$q_2 = 1 - \frac{4Ted}{(ed)^2} \int_0^\infty du (2 + Tedu)(1 + e^{-u})^{Ted} \frac{e^{-u}}{(1 + e^{-u})^2}. \quad (70)$$

The distribution function $P_2(q)$ remains non-trivial in finite dimension in the low-temperature phase

$$P_2(q) = T\gamma_{\min}\delta(q) + (1 - T\gamma_{\min})\delta(q - q_2) \quad (71)$$

where q_2 is given by (70) and γ_{\min} is

$$\gamma_{\min} = ed - 2 + \frac{1}{ed} \left[-\frac{19}{2} + 3g(Ted) - Ted g'(Ted) \right] + O(d^{-2}) \quad (72)$$

and where $g(t)$ is given by (42b).

So our $1/d$ expansion predicts a broken replica symmetry in the low-temperature phase in finite dimension.

One could extend the calculations of this section to overlaps between more than two walks and one would obtain expressions very similar to (70).

6. Generalisation to other distributions

In this section we discuss briefly the effect of using energy distributions other than (21).

In sections 3, 4 and 5 the high-dimension expansions were derived using the energy distribution (21), $\rho(\varepsilon) = \exp(-\varepsilon)$. This choice facilitated the calculation of the averages in table 1 (see table 2). It is possible to perform the same calculations using other distributions, $\rho(\varepsilon)$, for the disorder. However, for most choices, $\rho(\varepsilon)$, it is not possible to express the mean-field results (26)–(28) in a simple way in powers of d , and it is difficult to calculate the energy factors required for the diagrammatic expansion (table 1). To examine the effect of using different energy distributions, we shall therefore limit ourselves to

$$\rho(\varepsilon) = \frac{\varepsilon^{\alpha-1} \exp(-\varepsilon)}{\Gamma(\alpha)} \quad \alpha > 0, \varepsilon \geq 0. \quad (73)$$

This choice allows one to study changes produced by varying the parameter α , whilst still keeping the calculations relatively simple.

Let us first consider how the value of α affects the mean-field results. Using $\rho(\varepsilon)$ given by (73) with (24)–(26), one can show that, for large d , the mean-field energy in the low-temperature phase and transition temperature are given by

$$F_1(T) = \frac{\alpha}{ed^{1/\alpha}} + \frac{\alpha}{(ed^{1/\alpha})^2} + O(d^{-3/\alpha}) \quad T \leq T_c \quad (74)$$

and

$$T_c = \frac{1}{ed^{1/\alpha}} + \frac{2}{(ed^{1/\alpha})^2} + O(d^{-3/\alpha}). \quad (75)$$

One can see that, in general, we have an expansion in powers of $d^{1/\alpha}$ rather than d . However, when one moves away from the mean field, by evaluating the energy factors of table 1 for the diagrammatic expansion, one sees that these still behave as powers of d , not $d^{1/\alpha}$. For example the energy factor corresponding to figure 4(b) is given by

$$\begin{aligned} & \langle (\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T])^{\gamma T} - 2\langle \exp(-\gamma\varepsilon) \rangle^2 \\ &= -\frac{\Gamma(4\alpha)}{\alpha\Gamma^2(2\alpha)\gamma^{4\alpha}} + 2\frac{g_\alpha(t)}{\gamma^{4\alpha}} + O\left(\frac{1}{\gamma^{4\alpha+1}}\right) \end{aligned} \quad (76a)$$

where

$$g_\alpha(t) = \frac{t}{\Gamma^2(2\alpha)} \int_0^\infty \int_0^\infty dx dy x^{2\alpha-1} e^{-x} (x+ty)^{2\alpha-1} [(1+e^{-y})^t - 1]. \quad (76b)$$

So it still behaves as d^{-4} . When $\alpha = 1$, $g_\alpha(t) \equiv g(t)$ defined in (42b). Hence, for general α , one has an expansion in terms of $\gamma^{-\alpha}$ and γ^{-1} (or d^{-1} and $d^{-1/\alpha}$) to deal with. This makes calculations rather complicated and so we shall limit ourselves here to considering the leading behaviour of the specific heat.

The specific heat can still be found easily for arbitrary α . To do this in the low-temperature phase the behaviour of the function $g_\alpha(t)$ must be known for small t . This behaviour changes at $\alpha = \frac{1}{4}$. In the range $0 < \alpha \leq \frac{1}{4}$, $g_\alpha(t)$ behaves as $t^{4\alpha+1}$ for small t , whereas for $\alpha \geq \frac{1}{4}$ this changes to a t^2 dependence. This produces two possible behaviours for the low-temperature specific heat of the n -tree:

$$C_n = \begin{cases} \frac{\pi^2}{6} \frac{\Gamma(4\alpha-1)}{\Gamma^2(2\alpha)} \left(\frac{n-1}{n}\right) \frac{T}{(ed^{1/\alpha})^{2\alpha-1}} & \alpha \geq \frac{1}{4} \\ 4\alpha(4\alpha+1) \frac{J(\alpha)}{\Gamma^2(2\alpha)} \left(\frac{n-1}{n}\right) (ed^{1/\alpha})^{2\alpha} T^{4\alpha} & 0 < \alpha \leq \frac{1}{4} \end{cases} \quad (77a)$$

$$(77b)$$

where

$$J(\alpha) = \int_0^\infty \int_0^\infty dx dy x^{4\alpha-1} (y+y^2)^{2\alpha-1} \log(1+e^{-x}). \quad (77c)$$

Notice that for $\alpha \geq \frac{1}{4}$ one obtains a linear behaviour at low temperature, whilst for $0 < \alpha < \frac{1}{4}$ one has a $T^{4\alpha}$ behaviour. The low-temperature behaviour of the specific heat clearly depends on the nature of the disorder.

One should not be surprised by this change of behaviour for small values of α because one expects from (73) that the density of excitations around the ground state will increase as α becomes small.

7. Tests of the validity of the n -tree approximation method

The basic premise of the high-dimension expansion method, which has been presented in the preceding sections, is that results valid for the d -dimensional hypercubic lattice can be obtained by expanding the n -tree approximation for high dimension and then taking the limit $n \rightarrow \infty$. As was discussed in the closing paragraph of section 3, this procedure entails exchanging the limits $n \rightarrow \infty$ and $d \rightarrow \infty$. Although we have found that all quantities calculated so far have a finite limit for $n \rightarrow \infty$ when one uses this method, we have, as yet, no proof that these limits commute. In this section, therefore, we shall present a few problems for which the results can be tested (either against a solution obtained by a different approach, or against the results of simulations). First, we shall discuss the calculation, in high dimension, of the directed percolation threshold, p_c . Then we shall compare the results of the n -tree approximation with those of numerical simulation in dimension $d = 2$. Lastly we shall introduce the problem of directed polymers on a simplex and show that the n -tree approximation leads to the right answer for this problem.

7.1. The percolation threshold for d large

The first problem we shall consider, that of directed bond percolation, is simply the directed polymer problem with the random energies, ε_{ij} , chosen according to the distribution

$$\rho(\varepsilon) = p\delta(\varepsilon) + (1-p)\delta(\varepsilon-1) \quad (78)$$

with $0 \leq p \leq 1$. Again we shall look at the problem on a d -dimensional hypercubic lattice, directing the walks, which emanate from the origin, along the $(1, 1, \dots, 1)$ direction. If the weight p exceeds the percolation threshold, p_c , the ground-state energy per unit length is zero. The aim is to calculate this threshold, p_c , as a function of dimension, for d large.

Using the diagrammatic expansion method of section 3, one sees, from (40) and (41), that the ground-state energy per unit length, E_{GS} , of the n -tree approximation is given by

$$E_{GS} = -G_n(\gamma_{\min})|_{T=0} \quad (79a)$$

where

$$G_n(\gamma)|_{T=0} = \frac{1}{n\gamma} \log [d^n \langle \exp(-\gamma \varepsilon) \rangle^n + \frac{1}{2}(n-1)d^{n-1}(d-1) \langle \exp(-\gamma \varepsilon) \rangle^{n-2} \\ \times (\langle \max(\exp[-(\varepsilon_1 + \varepsilon_2)\gamma], \exp[-(\varepsilon_3 + \varepsilon_4)\gamma]) \rangle - 2\langle \exp(-\gamma \varepsilon) \rangle^2) + \dots] \quad (79b)$$

and γ_{\min} is defined as usual by (41c).

Using the percolation distribution (78), the averages can be performed to yield

$$G_n(\gamma)|_{T=0} = \frac{1}{n\gamma} \log [d^n [p + (1-p)e^{-\gamma}]^n + \frac{1}{2}(n-1)d^{n-1}(d-1)[p + (1-p)e^{-\gamma}]^{n-2} \\ \times \{2p^2 - p^4 + 4p(1-p)^2 e^{-\gamma} + (1-p)^4 e^{-2\gamma} \\ - 2[p + (1-p)e^{-\gamma}]^2\} + \dots]. \quad (80)$$

To obtain the percolation threshold, p_c , we require that when $p \geq p_c$ the ground-state energy is zero. So when $p \geq p_c$ the value, γ_{\min} , of γ that minimises $G_n(\gamma)$ is $\gamma_{\min} = \infty$. It is possible to show that $\gamma_{\min} = \infty$ as long as the argument of the logarithm in (80) is larger than 1 for $\gamma = \infty$. Therefore the percolation threshold is given by the condition that the argument of the logarithm is 1 for $\gamma = \infty$. This implies that

$$d^n p_c^n \left[1 - \left(\frac{n-1}{2} \right) \left(1 - \frac{1}{d} \right) p_c^2 + O(p_c^3) \right] = 1 \quad (81)$$

which becomes, when one includes higher-order diagrams,

$$\sum_{\text{diagrams}} A(\text{diagram}) B(\text{diagram}) = 1 \quad (82)$$

where the embedding counts, A , are the same as in table 1 and the factors B are now given for the percolation distribution (78) by table 4. As one needs here only $G_n(\gamma = \infty)$

Table 4. Energy factors, B , for the diagrams of figure 4 for the percolation distribution $\rho(\varepsilon) = p\delta(\varepsilon) + (1-p)\delta(\varepsilon-1)$ in the limit $\gamma = \infty$.

Diagram	Percolation factor
a	p^n
b	$-p^{n+2}$
c	$-p^{n+3}$
d	p^{n+4}
e	p^{n+4}
f	$-p^{n+4}$

for the distribution (78), the energy factors B depend only on p and are related to the percolation probability of each diagram. The solution, p_c , of (81) and (82) can be expanded in $1/d$, one gets

$$p_c = \frac{1}{d} + \left(\frac{n-1}{2n}\right) \frac{1}{d^3} + \left(\frac{2n-5}{2n}\right) \frac{1}{d^4} + \left(3 - \frac{99}{8n} + \frac{5}{8n^2}\right) \frac{1}{d^5} + O(d^{-6}). \quad (83)$$

Taking the limit $n \rightarrow \infty$, one recovers the results obtained by Blease (1977). So in this case the n -tree approximation method gives the right $1/d$ expansion.

7.2. The percolation problem in dimension $d = 2$

To further test the validity of the n -tree approximation method, we tried to calculate the ground-state energy per unit length, $E(p)$, for the distribution (78) in dimension two. A direct Monte Carlo simulation done on polymers of length 10^6 leads to $E(0.1) = 0.5537 \pm 0.0005$ and $E(0.2) = 0.3719 \pm 0.0005$.

By calculating the prediction, $E_n(p)$, of the n -tree approximation of this problem, we have obtained the results shown in figure 6 for $1 \leq n \leq 8$. When plotted against $n^{-1/3}$, the results converge linearly to the results of the simulation. So here again, the n -tree approximation seems, in the limit $n \rightarrow \infty$, to converge to the right answer. There is probably a way of relating the convergence rate to the known exponents $\omega = \frac{1}{3}$ and $\nu = \frac{2}{3}$, in dimension $d = 1 + 1$, but we were unable to find it.




Figure 6. The n -tree approximation for the ground-state energy of the percolation problem in dimension two. Approximations are shown for $p = 0.1$ (+) and $p = 0.2$ (x).

7.3. The simplex problem

Another test we considered for justifying the n -tree approximation was to study a simplified model of directed polymers in a random medium, the simplex model (see figure 7). The lattice in the transverse direction is a simplex and at each step along the preferred direction the polymer is allowed to jump from any corner of the simplex to any other corner.




Figure 7. The simplex model.

So in the simplex model the lattice consists of L planes with K points on each plane. Each point of a given plane is connected to all points of the previous and next plane. Therefore there are K^2 links between two consecutive planes. As always, on each link between two planes one chooses a random energy, ϵ , according to a given distribution $\rho(\epsilon)$. A directed walk is a walk which visits a single site in each plane, so the energy, E_w , of a walk of L steps is the sum of the energies of the bonds visited by the walk.

For this problem one can write the following recursion relation for the partition function:

$$Z_{L+1}(i) = \sum_{j=1}^K \exp[-\epsilon_{ij}(L)/T] Z_L(j) \quad (84)$$

where $Z_L(i)$ is the partition function of the walks which reach the i th point on plane L . So, the calculation of the free energy is the same as finding the largest Lyapounov exponent of a product of $K \times K$ matrices, the elements of which are random, positive, independent and identically distributed. To our knowledge, this is a problem which has not yet been solved for a general distribution $\rho(\epsilon)$.

We were only able to solve this problem in the particular case of zero temperature, large K , and the percolation distribution

$$\rho(\epsilon) = p\delta(\epsilon) + (1-p)\delta(1-\epsilon). \quad (85)$$

For large K it is convenient to use, instead of the parameter p (in (85)), the parameter x related to p by

$$p = K^{-x}. \quad (86)$$

Let us first show that the ground-state energy per unit length, E , is given by

$$E = \left(1 + \left[\frac{1}{x-1} \right] \right)^{-1} \quad x \geq 1 \quad (87)$$

where in (87) the square brackets denote the integer part of the argument. It is possible to derive this result by writing a Master equation for the probability, $P_L(E_1, \dots, E_i, \dots, E_K)$ that the ground-state energies of the walks of length L which reach the points $1, \dots, i, \dots, K$ of plane L are $E_1, \dots, E_i, \dots, E_K$. This Master equation can be solved using the fact that the differences in energy $|E_i - E_j|$ can only be 0 or 1, leading to (87) for large K . We shall only present here a simple argument which leads to (87).

It is easy to calculate the typical number of points in plane $L = m$ that are connected by a path of zero energy to the plane $L = 0$. This typical number is $K^{1+m(1-x)}$. As long as $1 + m(1-x)$ is positive, there is a path of zero energy from plane 0 to m . So if m is the largest integer such that

$$1 + m(1-x) > 0 \quad \text{and} \quad 1 + (m+1)(1-x) < 0 \quad (88)$$

the minimum energy to go from plane 0 to plane $m+1$ is 1. Therefore for large L the ground-state energy is $L/(m+1)$ where m satisfies (88), and so one gets (87).

Let us now present the n -tree approximation for this simplex problem. As in section 3, the ground-state energy E_n of the n th approximation will be given by

$$E_n = -\min_{\gamma} G_n(\gamma)|_{T=0}. \quad (89)$$

So we only need to determine the large- K behaviour of $G_n(\gamma)$. Clearly, from (13),

$$\begin{aligned} G_1(\gamma) &= \frac{1}{\gamma} \log\{K[K^{-x} + (1-K^{-x})e^{-\gamma}]\} \\ &\simeq \frac{1}{\gamma} \log(K^{1-x} + K e^{-\gamma}) \end{aligned} \quad (90)$$

so one has for large K

$$E_1 = \begin{cases} (x-1)/x & x \geq 1 \\ 0 & x \leq 1. \end{cases} \quad (91)$$

Going to the 2-tree approximation one can show that for large K

$$G_2(\gamma) \simeq \frac{1}{2\gamma} \log(K^{2-2x} + 2K^{2-x}e^{-\gamma} + K e^{-2\gamma}). \quad (92)$$

This can be understood because the average number of points which can be reached after two steps starting from a given point behaves like: K^{2-2x} for energy 0, K^{2-x} for energy 1 and K for energy 2. Minimising over γ (equation (89)), one finds that for large K

$$E_2 = \begin{cases} 0 & x \leq 1 \\ (x-1)/x & 1 \leq x \leq 2 \\ (2x-3)/2(x-1) & 2 \leq x. \end{cases} \quad (93)$$

Already with this 2-tree approximation, we see an improvement toward the true answer (87).

We did not find a simple way of solving the n -tree approximation for arbitrary x . However, for $x > 2$, we could generalise for arbitrary n the reasoning which led to the expression of $G_2(\gamma)$. For example, for odd n , one gets for $x > 2$

$$G_n(\gamma) \simeq \frac{1}{n\gamma} \log\left(\sum_{m=0}^{(n+1)/2} b_m K^{n-nx+mx} e^{-m\gamma} + \sum_{m=(n+3)/2}^n b_m K^{2n+1-2m-nx+mx} e^{-m\gamma}\right) \quad (94)$$

where b_m are constants independent of K .

This leads to

$$E_n = \begin{cases} 1 - 1/n(x-2) & \text{for } x > 2n/(n-1) \\ (x-1)/x & \text{for } 2n/(n-1) > x > 2 \end{cases} \quad (95)$$

which converges to the exact result $E = 1$ (see (87)) in the limit $n \rightarrow \infty$.

Similarly for even n and $x > 2$, one gets

$$E_n = \begin{cases} 1 - 1/n(x-2) & \text{for } x > (2n-2)/(n-2) \\ (2x-3)/2(x-1) & \text{for } (2n-2)/(n-2) > x > 2. \end{cases} \quad (96)$$

So with this example we see that the limits $K \rightarrow \infty$ and $n \rightarrow \infty$ commute at least for $x > 2$ and since K plays a similar role to d for the hypercubic lattice it supports the idea that these commute in general.

8. Conclusion

In this paper we have developed a method of obtaining $1/d$ expansions for the problem of directed polymers in a random medium, using the n -tree approximation. This consists of building a series of tree problems (n -trees) which give the hypercubic lattice in the limit $n \rightarrow \infty$. For our results to be valid one needs the limits $n \rightarrow \infty$ and $d \rightarrow \infty$ to commute. At present we have no proof of this, but in several examples, which can be solved independently (see section 7), we found that this n -tree method gave the correct answer.

We have obtained here the $1/d$ expansions of several quantities of interest for the directed polymer problem: free energy (42), specific heat (44), transition temperature (45), ground-state energy (46), transverse fluctuations (62) and overlaps (70). In contrast to the mean-field case, the low-temperature phase in finite dimension is no longer frozen. However, within our $1/d$ expansions, we find that the distribution of overlaps remains broad and that the transverse fluctuations remain linear in the length L of the polymer. This would be consistent with the idea that there exists a finite upper critical dimension above which the low-temperature phase has broken replica symmetry and the mean-field exponents.

It would be nice to see whether our results could be recovered by another approach which would not require the use of the n -tree approximation. It would also be interesting to know whether our $1/d$ expansions have a finite radius of convergence which could be identified with a finite upper critical dimension, or have a zero radius of convergence as has been noted for other $1/d$ expansions (Fisher and Singh 1989). This would require pushing our diagrammatic expansion further.

We think that the method presented here could be easily extended to other situations: finding the free energy as a function of the transverse displacement, site disorder rather than bond disorder. It should also be possible to extend it to other directed models and to growth models.

Lastly, in section 7, we introduced the simplex model which is, in spirit, in the directed polymer case, the equivalent of the Sherrington-Kirkpatrick (1975) model of spin glasses. This problem can be reduced to the calculation of the Lyapounov exponents for products of $K \times K$ random matrices, the elements of which are random, positive, independent and identically distributed. It would be nice to find a way of expanding the free energy of this model for large K .

Acknowledgments

We would like to thank H Spohn, J Vannimenus and J Yedidia for useful discussions. JC acknowledges support from SERC.

Appendix 1. Solution of the tree problem using replicas

In this appendix we derive the solution of the problem of directed polymers in a random medium on a tree using replicas. We also show how one can obtain the overlaps, q_2 , using this method, and derive (52b).

For generality, consider a tree with branching ratio K . Out of the K branches suppose that k_1 have random energies chosen according to the distribution $\rho_1(\epsilon)$, k_2

have random energies chosen according to $\rho_2(\varepsilon)$, etc... ($\sum_i k_i = K$). One can define a partition function for the problem, (2),

$$Z_L = \sum_w \exp(-E_w/T) \quad (\text{A1.1})$$

where the sum runs over all directed walks of length L and E_w is the total energy of the walk w (see (1)).

The free energy will be calculated using the replica trick (Sherrington and Kirkpatrick 1975)

$$\lim_{L \rightarrow \infty} \lim_{n \rightarrow 0} \frac{\log \langle Z_L^n \rangle}{nL} = \lim_{L \rightarrow \infty} \frac{\log Z_L}{L}. \quad (\text{A1.2})$$

$\langle Z_L^n \rangle$ can be thought of as the average of the partition function for n walks or replicas. We assume, following the Parisi scheme (Parisi 1980a, b) that the arrangements of the n walks which give the leading contribution to $\langle Z_L^n \rangle$ for large L are organised as follows:

- (i) the n walks remain together initially for a length Lq_1 ;
- (ii) the walks then split into m_1 groups of n/m_1 walks each and remain so for a length $L(q_2 - q_1)$;
- (iii) the walks then split into m_j groups of n/m_j walks each remaining so for a length $L(q_{j+1} - q_j)$;
- (iv) the walks finally separate into $m_M = n$ individual walks a length L from the starting point.

Therefore, one has

$$q_0 = 0 \leq q_1 \leq \dots \leq q_j \leq \dots \leq q_M = 1 \quad (\text{A1.3a})$$

$$1 = m_0 \leq m_1 \leq \dots \leq m_j \leq \dots \leq m_M = n. \quad (\text{A1.3b})$$

Then for large L , $\langle Z_L^n \rangle$ is dominated by

$$\langle (Z_L)^n \rangle \approx \max_{\{q_j\}} \max_{\{m_j\}} \prod_j \frac{m_j!}{\prod_i l_{ij}!} \prod_i k_i^{L(q_{j+1} - q_j)l_{ij}} \left(\int \rho_i(\varepsilon) \exp(-n\varepsilon/m_j T) d\varepsilon \right)^{L(q_{j+1} - q_j)l_{ij}} \quad (\text{A1.4})$$

where l_{ij} branches of class i are occupied by walks at the j th level and therefore $m_j = \sum_i l_{ij}$.

Taking the maximum over the l_{ij} , one gets

$$\langle (Z_L)^n \rangle \approx \max_{\{q_j\}} \max_{\{m_j\}} \prod_j \exp \left[L(q_{j+1} - q_j) m_j \log \left(\sum_i k_i \int \rho_i(\varepsilon) \exp(-n\varepsilon/m_j T) d\varepsilon \right) \right]. \quad (\text{A1.5})$$

To evaluate $\langle Z_L^n \rangle$ for large L one has to find the set $\{q_j\}$, $\{m_j\}$ which maximises the exponent in (A1.5). In the limit $n \rightarrow 0$, required in (A1.2), following Parisi (Parisi 1980a, b), we invert the inequality (A1.3b) to give

$$1 = m_0 \geq m_1 \geq \dots \geq m_j \geq \dots \geq m_M = n = 0 \quad (\text{A1.6})$$

and take the limit $M \rightarrow \infty$ by defining a function, $x(q_j)$, such that

$$m_j = \frac{n}{x(q_j)} \quad (\text{A1.7})$$

with q_j becoming a variable continuous on the interval $[0, 1]$ and $0 < x(q) < 1$. Then using (A1.2) and (A1.5) one obtains

$$\lim_{L \rightarrow \infty} \frac{1}{L} \log Z_L = \text{extr}_{x(q)} \int_0^1 \frac{dq}{x(q)} \log \left(\sum_i k_i \int \rho_i(\varepsilon) \exp(-\varepsilon x(q)/T) d\varepsilon \right). \quad (\text{A1.8})$$

As there is no explicit q dependence in (A1.8), $x(q)$ must be a constant. Hence one finds that

$$x(q)_{\text{optimum}} = \begin{cases} 1 & T \geq T_c \\ T\gamma_{\min} = T/T_c & T \leq T_c \end{cases} \quad (\text{A1.9})$$

where γ_{\min} is defined by

$$\left. \frac{dG(\gamma)}{d\gamma} \right|_{\gamma_{\min}} = 0 \quad (\text{A1.10})$$

with

$$G(\gamma) = \frac{1}{\gamma} \log \left(\sum_i k_i \int \rho_i(\varepsilon) e^{-\varepsilon\gamma} d\varepsilon \right). \quad (\text{A1.11})$$

(It is easy to show that $\gamma G(\gamma)$ is a convex function of γ , and one can therefore show that (A1.10) has a unique solution.) So one can conclude from (A1.8) and (A1.9) that

$$\lim_{L \rightarrow \infty} \frac{1}{L} \langle \log Z_L \rangle = \begin{cases} \frac{1}{T} G(1/T) & T \geq T_c \\ \frac{1}{T} G(\gamma_{\min}) & T \leq T_c. \end{cases} \quad (\text{A1.12})$$

This reduces to the result of section 2 (equations (13), (15)), derived using the travelling wave analogy, when one considers the case of a single distribution of disorder, $\rho(\varepsilon)$ (i.e. $k_1 = K$ and $\rho_1(\varepsilon) = \rho(\varepsilon)$). For clarity we shall now restrict ourselves to this problem.

From this replica calculation one can determine the overlaps. These are defined in section 2 (equations (17), (18)). The probability distribution of the overlaps, q_2 , between two walks can be calculated using the identification (Parisi 1983, Mezard *et al* 1984)

$$P_2(q) = \frac{dx(q)}{dq}. \quad (\text{A1.13})$$

Hence, from (A1.9), one can see that for $T \leq T_c$, $P_2(q)$ is composed of two delta functions

$$P_2(q) = T\gamma_{\min} \delta(q) + (1 - T\gamma_{\min}) \delta(q - 1) \quad T \leq T_c. \quad (\text{A1.14})$$

Above T_c , $P_2(q)$ is a single delta function at $q = 0$

$$P_2(q) = \delta(q) \quad T \geq T_c. \quad (\text{A1.15})$$

In the low-temperature phase $P_2(q)$ is not a single delta function, q_2 is not self-averaging and one can conclude that the phase has broken replica symmetry. This is analogous to the overlaps in the low-temperature phase of the mean-field spin glass (Mezard *et al* 1987).

One can, in addition, extract some information about the valley structure of the energy landscape from (A1.14). The replica symmetry breaking indicates a many-valley energy landscape. Assume one has a certain number of valleys, each valley α having a weight ω_α . If two walks fall in the same valley they have an overlap of unity and if they are in different valleys they have a zero overlap. Hence from (A1.14) it follows that

$$\sum_\alpha \omega_\alpha^2 = 1 - T\gamma_{\min} \quad T \leq T_c. \quad (\text{A1.16})$$

This is the result (52b) used in section 4.

Appendix 2. Derivation of equation (65)

The n -tree approximation to the d -dimensional lattice is a tree structure consisting of branches of a number of types. For example, the 2-tree consists of two types of branches, those which are formed by only one path and those which are made up of two paths (see figure 2(b)). In this appendix we show how one can calculate the fraction of each type of branch used by a typical walk on an n -tree. This quantity is needed in the calculation of the overlaps in section 5.

To see how one can evaluate these fractions, let us consider the simpler problem of directed walks on a tree when one has a discrete distribution of energies. Suppose that the probability of choosing a bond with energy ε_i is given by ρ_i , the ε_i forming a discrete set $\{\varepsilon_i\}$

$$\rho(\varepsilon) = \sum_i \rho_i \delta(\varepsilon - \varepsilon_i). \quad (\text{A2.1})$$

The free energy of the system per unit length, L , is then given by (15)

$$F(T) = \begin{cases} -G(\gamma_{\min}) & T \leq T_c \\ -G(1/T) & T \geq T_c \end{cases} \quad (\text{A2.2})$$

where

$$G(\gamma) = \frac{1}{\gamma} \log \left(\sum_i \rho_i \exp(-\gamma \varepsilon_i) \right) \quad (\text{A2.3})$$

and γ_{\min} is the unique solution of

$$\left. \frac{dG(\gamma)}{d\gamma} \right|_{\gamma_{\min}} = 0. \quad (\text{A2.4})$$

Let N_i denote the number of branches of type i in the walks. One can then write the partition function Z_L as

$$Z_L = \sum_{N_1=0}^L z_L(N_1) \exp(-\varepsilon_1 N_1 / T) \quad (\text{A2.5})$$

where $z_L(N_1)$ is the partial partition function for all walks of length L containing exactly N_1 steps on branches of type 1. Our aim is to calculate the average number of steps on branches of type 1

$$\langle \bar{N}_1 \rangle = \left\langle Z_L^{-1} \sum_{N_1=0}^L N_1 z_L(N_1) \exp(-\varepsilon_1 N_1 / T) \right\rangle \quad (\text{A2.6})$$

where $\bar{}$ denotes a thermal average. Comparing (A2.6) and (A2.5) we see that the average of N_1 can be obtained as a derivative of the free energy

$$\langle \bar{N}_1 \rangle = -T \frac{\partial}{\partial \varepsilon_1} \langle \log Z_L \rangle \quad (\text{A2.7})$$

and so, from (A2.2)–(A2.4) we see that

$$\langle \bar{N}_1 / L \rangle = \frac{\partial F}{\partial \varepsilon_1} = \frac{\rho_1 \exp(-\gamma \varepsilon_1)}{\sum_i \rho_i \exp(-\gamma \varepsilon_i)} \quad (\text{A2.8})$$

where $\gamma = \gamma_{\min}$ for $T \leq T_c$ and $\gamma = 1/T$ for $T \geq T_c$.

This idea can be extended to continuous distributions and used to calculate the average fraction of each type of branch used by walks on an n -tree. As the simplest example we shall take the 2-tree, although the extension to the n -tree is easy. From (32) and (36) the free energy per unit length of the 2-tree is given in the low-temperature phase by

$$F_2(T) = -\frac{1}{2\gamma_{\min}} \log[d(\exp(-\varepsilon\gamma_{\min}))^2 + \frac{1}{2}d(d-1) \times \langle \{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}} \rangle] \quad (\text{A2.9})$$

where γ_{\min} is defined by (32) and (36). To calculate the overlap for the 2-tree (see section 5) one needs to know the average fraction of branches used in a walk on the 2-tree which consist of two paths with energies $\varepsilon_1 + \varepsilon_2$ and $\varepsilon_3 + \varepsilon_4$ on the constituent links. Extending (A2.8) to continuous distributions and effective bonds, one can see from (A2.9) that this fraction is given in the low-temperature phase by

$$\frac{\frac{1}{2}d(d-1)(\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T])^{T\gamma_{\min}} \rho(\varepsilon_1)\rho(\varepsilon_2)\rho(\varepsilon_3)\rho(\varepsilon_4)}{d(\exp(-\varepsilon\gamma_{\min}))^2 + \frac{1}{2}d(d-1)\langle \{\exp[-(\varepsilon_1 + \varepsilon_2)/T] + \exp[-(\varepsilon_3 + \varepsilon_4)/T]\}^{T\gamma_{\min}} \rangle}. \quad (\text{A2.10})$$

This is the result quoted in (65). This method can easily be extended to calculate the average fraction of its length spent by a walk on any type of branch on the n -tree.

References

- Bleale J 1977 *J. Phys. C: Solid State Phys.* **10** 925
 Bramson M 1983 Convergence of solutions of the Kolmogorov equation to travelling waves *Memories of the American Mathematical Society* **285**
 Bray A J and Moore M A 1979 *J. Phys. C: Solid State Phys.* **12** 79
 Cook J and Derrida B 1989a *Europhys. Lett.* **10** 195
 — 1989b *J. Stat. Phys.* **57** 89
 de Dominicis C and Kondor I 1985 *J. Physique Lett.* **46** L1037
 Derrida B and Gardner E 1986 *J. Phys. C: Solid State Phys.* **19** 5783
 Derrida B and Spohn H 1988 *J. Stat. Phys.* **51** 817
 Dhar D 1988 *Phys. Lett.* **130A** 308
 Fisher M E and Singh R R P 1989 *Disorder in Physical Systems* ed G Grimmett and D J A Welsh, to appear
 Forster D, Nelson D R and Stephen M J 1977 *Phys. Rev. A* **16** 732
 Halpin-Healy T 1989 *Phys. Rev. Lett.* **62** 442
 Hammersley J M 1974 *Ann. Prob.* **2** 652
 Huse D A and Fisher D S 1987 *J. Phys. A: Math. Gen.* **20** L997
 — 1988 *Phys. Rev. B* **38** 386
 Huse D A, Henley C L and Fisher D S 1985 *Phys. Rev. Lett.* **55** 2924
 Imbrie J Z and Spencer T 1988 *J. Stat. Phys.* **52** 609
 Kardar M 1987 *J. Appl. Phys.* **61** 3601
 Kardar M, Parisi G and Zhang Y C 1986 *Phys. Rev. Lett.* **56** 889
 Kardar M and Zhang Y C 1987 *Phys. Rev. Lett.* **58** 2087
 Kim J M and Kosterlitz J M 1989 *Phys. Rev. Lett.* **62** 2289
 Krug J 1987 *Phys. Rev. A* **36** 5465
 Krug J and Spohn H 1988 *Phys. Rev. A* **38** 4271
 McKane A J and Moore M A 1988 *Phys. Rev. Lett.* **60** 527
 Mezard M, Parisi G, Sourlas N, Toulouse G and Virasoro M 1984 *J. Physique* **45** 843
 Mezard M, Parisi G and Virasoro M 1987 *Spin Glass Theory and Beyond* (Singapore: World Scientific)
 Nattermann T and Renz W 1988 *Phys. Rev. B* **38** 5184

Parisi G 1980a *J. Phys. A: Math. Gen.* **13** 1101

— 1980b *J. Phys. A: Math. Gen.* **13** 1887

— 1983 *Phys. Rev. Lett.* **50** 1946

Renz W 1989 to be published

Sherrington D and Kirkpatrick S 1975 *Phys. Rev. Lett.* **32** 1792

Sourlas N 1986 *Europhys. Lett.* **1** 189

Wolf D E and Kertész J 1987 *Europhys. Lett.* **4** 651

Zhang Y C 1987 *Phys. Rev. Lett.* **59** 2125