

Tutorial 12, Statistical Mechanics: Concepts and applications 2016/17 ICFP Master (first year)

Maurizio Fagotti, Olga Petrova, Werner Krauth
Tutorial exercises

I. WORKSHEET: FIRST-ORDER PHASE TRANSITIONS




FIG. 1. Left: The six possible orientations of a molecule. Right: The matrix representation of the orientations.

Source: R. A. Sauerwein and M. J. De Oliveira, *J. of Chem. Phys.* **144**, 194904 (2016).

1. Lattice model for biaxial and uniaxial liquid crystals

Reminder: Liquid crystals are states of matter that can be thought of as something in between conventional liquids and solid crystals. They can exhibit various phases, one of the most common being the nematic phase. In nematics, the molecules tend to have the same alignment but their positions are not correlated. Thus, their center of mass positions are randomly distributed as in a liquid, but still maintain a long-range directional order.

The molecules are generally three-dimensional objects, often elongated in one direction. If there is a preferred direction for one of the axis (most commonly, the major one), the nematic is said to be *uniaxial*. If there is a secondary preferred direction that is perpendicular to the other one, such a nematic is called *biaxial*. At sufficiently high temperature, thermal fluctuations are able to destroy the ordering of the nematic phase, and the system reaches a conventional isotropic liquid phase.




FIG. 2. Sketch of the phase diagram in the variables h (cf. (2)) and $\Theta = \frac{12kT}{J\gamma r^2(h^2+3)}$, where γ is the coordination number. The phases are isotropic (I) uniaxial nematic (N_{\pm}), and biaxial nematic (B).

The model: Although we emphasize that there is no positional order (i.e., underlying lattice structure) in liquid crystals, we start with a simple lattice model in order to describe the phase transitions in nematics. In particular, we consider a regular lattice where, at each site, there is a parallelepiped-shaped molecule. We assume that the molecules can take only the 6 orientations shown in Fig. 1.

A rod-like nematic phase occurs when the major axes are mostly parallel to each other whereas the other two axes point in arbitrary directions. A disc-like nematic phase occurs when the minor axes are mostly parallel to each other whereas the other two axes point in arbitrary direction. A biaxial phase takes place when both the major axes are mostly parallel to each other, and the minor axes are mostly parallel to each other. In this case, same can be said about the third axis as well.

The system is described by the Hamiltonian

$$H = -J \sum_{(i,j)} \text{tr}[\Omega^i \Omega^j], \quad (1)$$

where the sum is over nearest neighbor pairs, $\Omega_i \in \{W_1, \dots, W_6\}$ are the diagonal matrices shown in fig. 1 (for the moment, you can interpret a , b , and c as the lengths in the various directions).

- (a) Show that, without loss of generality, the matrices W_i can be chosen to be traceless, i.e. $a+b+c=0$.
- (b) Within the mean-field approximation, neglect the fluctuations, i.e. use the following approximation: $(x - \langle x \rangle)(y - \langle y \rangle) \rightarrow 0$. Define $Q_{\mu} = \langle \Omega_{\mu\mu}^i \rangle$ and write down an effective mean-field Hamiltonian linear in Ω^i . The coordination number of the lattice is z .
- (c) Within mean field, calculate the free energy and write down the system of equations satisfied by the diagonal elements of Q . The total number of molecules is N , and you can leave the equations in matrix form.

(d) Use the following parametrization

$$Q = \begin{pmatrix} -\frac{q+\eta}{2} & 0 & 0 \\ 0 & -\frac{q-\eta}{2} & 0 \\ 0 & 0 & q \end{pmatrix} \quad W_1 = \begin{pmatrix} -\frac{\tau}{6}(h-3) & 0 & 0 \\ 0 & -\frac{\tau}{6}(h+3) & 0 \\ 0 & 0 & \frac{\tau}{3}h \end{pmatrix}. \quad (2)$$

Identify the phases of matter parametrized by q, η .

- (e) Write down the free energy in the uniaxial (rod-like and disk-like) nematic phases. Identify (i.e., write down the equation for) the line of the phase transition from the isotropic phase to the nematic phase. What can be an order parameter for the phase transition? Is it a first-order or a continuous transition (is there a jump in the order parameter)?
- (f) The uniaxial-biaxial nematic transition is continuous. Identify the line of the phase transition. A Landau point is an isolated critical point on a line of first-order transitions. Here it occurs at $\Theta = 1, h = 0$. The order parameter is small in the vicinity of the Landau point, so the free energy can be expanded in powers of the elements of Q .
- (g) Determine the line of the isotropic to uniaxial nematic transition close to the Landau point. Calculate the jump in the order parameter q .